



Mellin Transforms and Asymptotics: Harmonic Sums

Philippe Flajolet, Xavier Gourdon, Philippe Dumas

► To cite this version:

Philippe Flajolet, Xavier Gourdon, Philippe Dumas. Mellin Transforms and Asymptotics: Harmonic Sums. [Research Report] RR-2369, INRIA. 1994. inria-00074307

HAL Id: inria-00074307

<https://inria.hal.science/inria-00074307>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

Mellin Transforms and Asymptotics: Harmonic Sums

Philippe FLAJOLET - Xavier GOURDON
Philippe DUMAS

N° 2369

Octobre 1994

PROGRAMME 2



Les rapports de recherche de l'INRIA
sont disponibles en format postscript sous
ftp.inria.fr (192.93.2.54)

si vous n'avez pas d'accès ftp
la forme papier peut être commandée par mail :
e-mail : dif.gesdif@inria.fr
(n'oubliez pas de mentionner votre adresse postale).

par courrier :
Centre de Diffusion
INRIA
BP 105 - 78153 Le Chesnay Cedex (FRANCE)

INRIA research reports
are available in postscript format
ftp.inria.fr (192.93.2.54)

if you haven't access by ftp
we recommend ordering them by e-mail :
e-mail : dif.gesdif@inria.fr
(don't forget to mention your postal address).

by mail :
Centre de Diffusion
INRIA
BP 105 - 78153 Le Chesnay Cedex (FRANCE)

Mellin Transforms and Asymptotics: Harmonic Sums

Philippe FLAJOLET, Xavier GOURDON, and Philippe DUMAS
Algorithms Project, INRIA Rocquencourt

Abstract. *This survey presents a unified and essentially self-contained approach to the asymptotic analysis of a large class of sums that arise in combinatorial mathematics, discrete probabilistic models, and the average-case analysis of algorithms. It relies on the Mellin transform, a close relative of the integral transforms of Laplace and Fourier. The method applies to harmonic sums that are superpositions of rather arbitrary “harmonics” of a common base function. Its principle is a precise correspondence between individual terms in the asymptotic expansion of an original function and singularities of the transformed function. The main applications are in the area of digital data structures, probabilistic algorithms, and communication theory.*

Transformation de Mellin et asymptotique: Sommes harmoniques

Résumé. Cet article de synthèse présente une approche unifiée à l'analyse asymptotique d'une large classe de sommes qui se présentent en analyse combinatoire, dans l'étude des modèles probabilistes discrets ainsi qu'en analyse d'algorithmes. L'approche repose sur la transformation de Mellin qui est étroitement liée aux transformations intégrales de Laplace et Fourier. Elle s'applique aux sommes harmoniques, lesquelles sont des superpositions linéaires d'“harmoniques” d'une fonction de base commune. Le principe en est une correspondance précise entre les termes du développement asymptotique de la fonction d'origine et les singularités de sa transformée. Les principales applications se situent dans le domaine des structures de donnée digitales, des algorithmes probabilistes, et de la théorie de la communication.

MELLIN TRANSFORMS AND ASYMPTOTICS: HARMONIC SUMS

PHILIPPE FLAJOLET, XAVIER GOURDON, AND PHILIPPE DUMAS

*Dedicated to Don Knuth and N. G. de Bruijn
for their pioneering works on Mellin transforms and combinatorics.*

ABSTRACT. This survey presents a unified and essentially self-contained approach to the asymptotic analysis of a large class of sums that arise in combinatorial mathematics, discrete probabilistic models, and the average-case analysis of algorithms. It relies on the Mellin transform, a close relative of the integral transforms of Laplace and Fourier. The method applies to harmonic sums that are superpositions of rather arbitrary “harmonics” of a common base function. Its principle is a precise correspondence between individual terms in the asymptotic expansion of an original function and singularities of the transformed function. The main applications are in the area of digital data structures, probabilistic algorithms, and communication theory.

*Die Theorie der reziproken Funktionen und Integrale
ist ein centrales Gebiet, welches manche anderen Gebiete
der Analysis miteinander verbindet.*

— HJALMAR MELLIN

INTRODUCTION

Hjalmar Mellin (1854–1933, see [59] for a summary of his works) gave his name to the *Mellin transform* that associates to a function $f(x)$ defined over the positive reals the complex function $f^*(s)$ where

$$(1) \quad f^*(s) = \int_0^\infty f(x)x^{s-1} dx.$$

The change of variables $x = e^{-u}$ shows that the Mellin transform is closely related to the Laplace transform and the Fourier transform. However, despite this connexion, there are numerous applications where it proves convenient to operate directly

with the Mellin form (1) rather than the Laplace-Fourier version. This is often the case in complex function theory (asymptotics of Gamma-related functions [64, 65]), in number theory (coefficients of Dirichlet series, after Riemann), in applied mathematics (asymptotic estimation of integral forms), and in the analysis of algorithms (harmonic sums introduced below). Thus, throughout this paper, we operate directly with the Mellin transform.

The major use of Mellin transform examined here is for the *asymptotic analysis* of sums obeying the general pattern

$$(2) \quad G(x) = \sum_k \lambda_k g(\mu_k x),$$

either as $x \rightarrow 0$ or as $x \rightarrow \infty$. Following a proposal of [31], sums of this type are called *harmonic sums* as they represent a superposition of “harmonics” of a single base function $g(x)$.

Harmonic sums surface at many places in combinatorial mathematics as well as in the analysis of algorithms and data structures. De Bruijn and Knuth are responsible in an essential way for introducing the Mellin transform in this range of problems, as attested by Knuth’s account in [56, p. 131] and the classic paper [16] which have been the basis of many later combinatorial applications. For instance, the analyses of the radix-exchange sort algorithm [56] and of the expected height of plane (Catalan) trees [16] involve the quantities

$$(3) \quad S_n = \sum_{k=0}^{\infty} \left[1 - \left(1 - \frac{1}{2^k} \right)^n \right] \quad \text{and} \quad T_n = \sum_{k=0}^n d(k) \frac{\binom{2n}{n-k}}{\binom{2n}{n}},$$

where $d(k)$ is the number of divisors of k and $\binom{a}{b}$ is the binomial coefficient. These discrete quantities have continuous analogues that are harmonic sums

$$(4) \quad G(x) = \sum_{k=0}^{\infty} \left[1 - e^{-x/2^k} \right] \quad \text{and} \quad H(x) = \sum_{k=0}^{\infty} d(k) e^{-k^2 x^2},$$

and elementary arguments establish that asymptotically $S_n \sim G(n)$ and $T_n \sim H(n)$.

There are inherent difficulties in the asymptotic analysis of the discrete sums S_n, T_n or their continuous analogues $G(x), H(x)$. The divisor function in T_n and $H(x)$ fluctuates heavily in a rather irregular manner (for instance it equals 2 if and only if k is prime). The quantities S_n and $G(x)$ look more innocuous; however, a Mellin based analysis to be shown later reveals that they involve subtle fluctuations of a tiny amplitude, less than 10^{-5} . Such behaviours preclude the use of elementary real asymptotic techniques in most sums of this type.

Over the past twenty years, perhaps some fifty odd analyses of algorithms or related evaluations of parameters of combinatorial structures have involved in a crucial way a Mellin treatment of harmonic sums of sorts. We shall organize and detail in later sections several of these examples that can be broadly categorized, in terms of their range of applications, as follows.

- Digital searching methods: radix-exchange sort, digital trees, digital search trees [56] and their generalizations like “bucket” trees [32], Patricia trees [56, 80], suffix trees [48].
- Digital trees, variance and biased-bit models [54, 79].
- Tree manipulation algorithms: height and stack depth [16], register allocation [50].
- String searching and the occurrence of patterns in strings [68] (and references therein), with applications to carry propagation in binary adders [57] and data compression [48, 47].
- Multidimensional searching problems [29] and Euclidean matching [74].
- Extendible hashing and grid file methods [72, 73].
- Parallel and distributed algorithms like the leader election technique of [70], communication protocols based on the Capetanakis–Tsybakov scheme [20, 39, 46, 63], parallel sorting networks based on Batcher’s odd-even scheme [75].
- Divide-and-conquer algorithms: mergesort [24] and geometric maxima finding [24].
- Randomized data structures, like skip lists [52].
- Probabilistic estimation algorithms: probabilistic counting [27], adaptive sampling [23], or approximate counting [22, 71].

In many cases, notably digital search trees, harmonic sums are an alternative to the method of “Rice integrals” that is discussed in detail elsewhere [34].

Some brief accounts of the method are given in the books by Hofri [44, p. 48ff], Kemp [51, p. 141], Mahmoud [62], and in the handbook chapter [84]. We follow here the architecture of the informal survey [31].

General properties of the Mellin transform are usually treated in detail in books on integral transforms, like those of Doetsch [18], Widder [86], or Titchmarsh [82]. Asymptotic methods in connection with Mellin transforms are discussed within the context of applied mathematics in treatises by Davies [13], Dingle [17], and Wong [88]. In particular, our work is close in spirit to Wong’s who discusses extensively an analogue of harmonic sums in the form of “harmonic integrals”

$$(5) \quad G(x) = \int_0^\infty \lambda(\kappa) g(\mu(\kappa)x) d\kappa.$$

(In this continuous case, by a change of variables, one may always assume that $\mu(k) \equiv k$.) Some of the many uses in number theory are treated for instance in [4, 7, 42].

Principles. A major use of Mellin transforms in asymptotic analysis is for estimating asymptotically harmonic sums (2). The λ_k are the “amplitudes”, the μ_k are the “frequencies”, and $g(x)$ is the “base function”. Harmonic sums reduce to usual Fourier series when the base function $g(x)$ is taken to be a complex exponential, $g(x) = e^{\pm ix}$, and the frequencies are the integers, $\mu_k \equiv k$.

1. *Mellin transforms and the “separation” property.* The Mellin transform as defined in (1) converges in a strip of the complex plane called the fundamental

strip. By a direct change of variables, the Mellin transform of $g(\mu x)$ is μ^{-s} times the transform $g^*(s)$ of $g(x)$. Thus, by linearity, the Mellin transform of a general harmonic sum is (conditionally)

$$(6) \quad G^*(s) = \Lambda(s) \cdot g^*(s),$$

where

$$\Lambda(s) = \sum_k \lambda_k \mu_k^{-s}, \quad g^*(s) = \int_0^\infty g(x) x^{s-1} dx.$$

It therefore factors as the product of the transform of the base function and of a generalized Dirichlet series: *Mellin transforms applied to harmonic sums "separate" the amplitude-frequency pair from the base function.*

The inversion theorem for Mellin transforms is analogous to Fourier inversion,

$$(7) \quad f(x) = \frac{1}{2i\pi} \int_{c-i\infty}^{c+i\infty} f^*(s) x^{-s} ds.$$

There, the integration line $\Re(s) = c$ should be taken in the fundamental strip of the Mellin transform.

Basic functional properties of the Mellin transform are recalled in Section 1 (Theorem 1) and the separation property is expressed by Lemma 2 of Section 3.

2. Poles of transforms and asymptotics of originals. There is a fundamental correspondence between the asymptotic expansion of a function (either at 0 or ∞) and singularities of the transformed function.

First and foremost, Mellin asymptotics crucially relies on *analytic continuation* of transforms. Consider the problem of asymptotically expanding $f(x)$ as $x \rightarrow 0$ when $f^*(s)$ is known to be meromorphically continuable in \mathbb{C} . The starting point is the inversion formula (7). The line of integration is then shifted to the *left* while taking residues into account. For instance when $c > 0$ and $f^*(s)$ has simple poles at the nonpositive integers, each simple pole at $s = -m$ contributes a term proportional to x^m since

$$\text{Res}(f^*(s)x^{-s})_{s=-m} = \text{Res}(f^*(s))_{s=-m} \cdot x^m.$$

(There $\text{Res}(g(s))_{s=s_0}$ denotes the residue of $g(s)$ at $s = s_0$.) Thus, globally by integrating along an infinite rectangle with sides $\Re(s) = c$ and $\Re(s) = -M - 1/2$ (M an integer) gives by the residue theorem

$$(8) \quad f(x) = \sum_{m=0}^M (\text{Res} f^*(s))_{s=-m} x^m + \mathcal{O}(x^{M+1/2}).$$

Some additional growth condition on $f^*(s)$ is evidently necessary in (8).

The computation outlined above reflects a general phenomenon: *Poles of a Mellin transform are in direct correspondence with terms in the asymptotic expansion of the original function at either 0 or $+\infty$.*

For the asymptotic evaluation of a harmonic sum $G(x)$ this principle applies to $G^*(s)$ provided the Dirichlet series $\Lambda(s)$ and the transform $g^*(s)$ are each analytically continuable and of controlled growth. This assumption is realistic for

many naturally occurring base functions and for amplitude and frequencies given by “laws” that are regular enough.

The fundamental correspondence is explored in Section 2 (Theorems 3, 4); its application to harmonic sums is spelled out in Section 3.

Example. The mode of operation of Mellin asymptotics is well illustrated by the harmonic sum

$$(9) \quad G(x) = \sum_{k=1}^{\infty} (-1)^k (\log k) e^{-k^2 x}.$$

In this case, the classical equations

$$\Gamma(s) = \int_0^{\infty} e^{-x} x^{s-1} dx \quad \text{and} \quad \zeta(s) = \sum_{k=1}^{\infty} \frac{1}{k^s},$$

define the Gamma function of Euler and the Riemann zeta function [85]. Thus, $\Gamma(s)$ is the Mellin transform of the base function $g(x) = e^{-x}$ and a simple computation yields

$$\Lambda\left(\frac{s}{2}\right) = \frac{d}{ds} (1 - 2^{1-s}) \zeta(s).$$

In this way, the transform of $G(x)$ is found to be

$$(10) \quad G^*(s) = [2^{1-2s} (\log 2) \zeta(2s) + (1 - 2^{1-2s}) \zeta'(2s)] \cdot \Gamma(s).$$

Analytically, the representation (10) is easily justified when $\Re(s) > 1$, a condition that ensures simultaneously absolute convergence of the Dirichlet series and of the Mellin integral.

It is known that the Gamma function and the zeta function are both continuable to the complex plane: $\Gamma(s)$ has simple poles at the nonpositive integers while $\zeta(s)$ has only a simple pole at $s = 1$. Also, $\Gamma(s)$ decreases fast along vertical lines while $\zeta(s)$ is only of moderate growth. Thus, globally, $G^*(s)$ gets small for $s \rightarrow \pm i\infty$ and the integration contour can legitimately be shifted to the left. For instance, by sweeping the integration contour till the vertical line $\Re(s) = -5/2$, one gets from the poles at $s = 0, -1, -2$

$$G(x) = \log \sqrt{\frac{\pi}{2}} + c_1 x + c_2 x^2 + \mathcal{O}(x^{5/2}),$$

with $c_1 = 7\zeta'(-2)$ and $c_2 = -\frac{31}{2}\zeta'(-4)$. The first few terms are then determined in a matter of seconds with the help of a computer algebra system like Maple that “knows” the expansions of $\zeta(s)$ and $\Gamma(s)$ at points of interest, like $\zeta(0) = -\frac{1}{2}$, $\zeta'(0) = -\log \sqrt{2\pi}$.

The computation is easily carried out to any order, and one finds

$$G(x) \sim \log \sqrt{\frac{\pi}{2}} + \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k!} (2^{1+2k} - 1) \zeta'(-2k) x^k.$$

Such an expansion might be obtained otherwise via a reduction to Euler-Maclaurin summation but application of this formula would be hampered by the alternation of sign and by the presence of the logarithmic factor. Mellin asymptotics is the approach of choice here.

The method is susceptible to a large number of variations. It is applicable each time the Dirichlet series of an amplitude-frequency pair is reducible to a standard function, like for instance

$$\sum_{k=0}^{\infty} \frac{1}{(2^k)^s} = \frac{1}{1-2^{-s}}, \quad \sum_{k=1}^{\infty} \frac{d(k)}{k^s} = \zeta^2(s).$$

In this way sums involving geometrically increasing frequencies or highly fluctuating amplitudes of an arithmetic nature can be analyzed effortlessly, see Sections 4–7. Another feature is that the correspondence between poles of transforms, and asymptotic terms of originals fares both ways; this fact allows for base functions that admit of no explicit transform, like

$$g(x) = e^{-x^2} \sqrt{1+x},$$

as well as exotic Dirichlet series like

$$\Lambda(s) = \sum_{k=1}^{\infty} \frac{\log k}{\sqrt{1+k^2}} \frac{(-1)^k}{k^s}.$$

This flexibility explains the power of the Mellin method in asymptotic analysis, which would apply for instance to a harmonic sum like

$$G(x) = \sum_{k=1}^{\infty} (-1)^k (\log k) e^{-k^2 x^2} \sqrt{\frac{1+kx}{1+k^2}}.$$

The general methodology for analyzing such “implicit” sums is discussed in Section 8.

Plan of the paper. Part I is devoted to the general functional and asymptotic properties of Mellin transforms. Our presentation assumes only rudiments of complex analysis (contour integrals, residue theorem) as found for instance in [81, 85]. The general framework is conveniently built on the Lebesgue integral [81, Ch. X] allowing dominated convergence properties rather than the Riemann integral. This distinction is however somewhat immaterial as specific examples can be dealt with using Riemann integrals only. A number of examples related to combinatorial analysis and the analysis of algorithms are then presented in Part II. Finally, methods for dealing with wider classes of sums form the subject of Part III.

I. Mellin Transforms and Asymptotics

In this part, we lay down the basic framework of Mellin asymptotics. Section 1 gives the basic functional properties of Mellin transforms and Section 2 discusses the fundamental correspondence between asymptotic properties of an original function and singularities of its transform. Section 3 develops the basic treatment of

harmonic sums that form the subject of this paper. Section 4 briefly examines consequences for general summatory formulæ.

1. BASIC PROPERTIES

We start by recalling the salient properties of the Mellin transform. The definition domain of a Mellin transform turns out to be a strip. We thus introduce the notation $\langle \alpha, \beta \rangle$ for the *open strip* of complex numbers $s = \sigma + it$ such that $\alpha < \sigma < \beta$.

Definition 1. Let $f(x)$ be locally Lebesgue integrable over $(0, +\infty)$. The Mellin transform of $f(x)$ is defined by

$$(11) \quad \mathcal{M}[f(x); s] = f^*(s) = \int_0^{+\infty} f(x)x^{s-1} dx,$$

The largest open strip $\langle \alpha, \beta \rangle$ in which the integral converges is called the *fundamental strip*.

From the decomposition $\int_0^\infty = \int_0^1 + \int_1^\infty$, we see that α is the infimum of all A such that $f(x)x^{A-1}$ is integrable over $(0, 1]$ and β is the supremum of all B such that $f(x)x^{B-1}$ is integrable over $[1, +\infty)$. Most functions have an order at 0 and ∞ so that an existence strip for $f^*(s)$ can be guaranteed.

Lemma 1. The conditions

$$f(x) \underset{x \rightarrow 0^+}{=} \mathcal{O}(x^u), \quad f(x) \underset{x \rightarrow +\infty}{=} \mathcal{O}(x^v),$$

when $u > v$, guarantee that $f^*(s)$ exists in the strip $\langle -u, -v \rangle$.

Thus, existence of a Mellin transform is granted for locally integrable functions¹ whose exponent in the order at 0 is strictly larger than the exponent of the order at infinity. The asymptotic form of $f(x)$ at 0 constrains the leftmost boundary of the fundamental strip of $f^*(s)$; the asymptotic form at $+\infty$ constrains the rightmost boundary. Monomials x^c , including constants, thus do not have transforms under Definition 1.

As the integral defining $f^*(s)$ depends analytically on the complex parameter s , a Mellin transform is in addition analytic in its fundamental strip.

For instance, the function $f(x) = (1+x)^{-1}$ is $\mathcal{O}(x^0)$ at 0 and $\mathcal{O}(x^{-1})$ at infinity, hence a guaranteed existence strip for $f^*(s)$ that is $\langle 0, 1 \rangle$, which here coincides with the fundamental strip. In this case, the Mellin transform may be found from the classical Beta integral [85, p. 254] to be

$$f^*(s) = \frac{\pi}{\sin \pi s},$$

which is analytic in $\langle 0, 1 \rangle$, as predicted.

The function $g(x) = e^{-x}$ satisfies

$$e^{-x} \underset{x \rightarrow 0^+}{\sim} 1, \quad e^{-x} \underset{x \rightarrow +\infty}{=} \mathcal{O}(x^{-b}) \quad \text{for any } b > 0$$

¹In the sequel, all functions considered are tacitly assumed to be locally integrable.

so that its transform, known as the Gamma function [85, Ch. XII]

$$(12) \quad \Gamma(s) = \int_0^{+\infty} e^{-x} x^{s-1} dx,$$

is *a priori* defined in $\langle 0, +\infty \rangle$ and analytic there.

Let $H(x)$ be the (Heaviside-like) step function defined by

$$(13) \quad H(x) = 1 \quad \text{if } 0 \leq x < 1, \quad H(x) = 0 \quad \text{if } 1 < x.$$

Then,

$$H^*(s) = \frac{1}{s} \quad s \in \langle 0, +\infty \rangle.$$

The complementary function $\overline{H}(x) = 1 - H(x)$ satisfies $\overline{H}^*(s) = -s^{-1}$ for $s \in \langle -\infty, 0 \rangle$, where the fundamental strips corresponding to H and \overline{H} are disjoint.

Functional properties. Simple changes of variables in the definition of Mellin transforms yield a basic set of transformation rules summarized in Fig. 1 and Theorem 1 below.

Theorem 1 (Functional properties). *Let $f(x)$ be a function whose transform admits the fundamental strip $\langle \alpha, \beta \rangle$. Let ρ be a nonzero real number, and μ, ν be positive reals. Then the following relations hold:*

$$\begin{aligned} \mathcal{M}[f(\mu x); s] &= \mu^{-s} f^*(s) & s \in \langle \alpha, \beta \rangle \\ \mathcal{M}[\sum_k \lambda_k f(\mu_k x); s] &= (\sum_{k \in \mathcal{K}} \lambda_k \mu_k^{-s}) \cdot f^*(s) & \mathcal{K} \text{ finite, } \lambda_k > 0 \\ \mathcal{M}[x^\nu f(x); s] &= f^*(s + \nu) & s \in \langle \alpha, \beta \rangle \\ \mathcal{M}[f(x^\rho); s] &= \frac{1}{\rho} f^*\left(\frac{s}{\rho}\right) & s \in \langle \rho\alpha, \rho\beta \rangle. \end{aligned}$$

The most important rule is the *rescaling rule* that gives the transform of $f(\mu x)$ as $\mu^{-s} f^*(s)$ via the change of variables $x \mapsto \mu x$, provided $\mu > 0$. By linearity of the transform, one also has

$$\mathcal{M}\left[\sum_k \lambda_k f(\mu_k x); s\right] = \left(\sum_k \frac{\lambda_k}{\mu_k^s}\right) \cdot f^*(s),$$

whenever k ranges over a finite set of indices. This formula can usually be extended to infinite sums that define the harmonic sums already mentioned in the introduction, see Lemma 1 and Theorem 5 below. For instance from

$$f(x) = \frac{e^{-x}}{1 - e^{-x}} = e^{-x} + e^{-2x} + e^{-3x} + \dots,$$

one finds

$$f^*(s) = \Gamma(s) \cdot \left(\frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \dots\right) = \Gamma(s)\zeta(s),$$

as long as $\Re(s) > 1$, a condition that simultaneously entails absolute convergence of the Mellin integral of $f(x)$ and of the sum defining the zeta function.

	$f(x)$	$f^*(s)$	$\langle \alpha, \beta \rangle$	
F_1	$x^\nu f(x)$	$f^*(s + \nu)$	$\langle \alpha - \nu, \beta - \nu \rangle$	
F_2	$f(x^\rho)$	$\frac{1}{\rho} f^*\left(\frac{s}{\rho}\right)$	$\langle \rho\alpha, \rho\beta \rangle$	$\rho > 0$
	$f(1/x)$	$f^*(-s)$	$\langle -\beta, -\alpha \rangle$	
F_3	$f(\mu x)$	$\frac{1}{\mu^s} f^*(s)$	$\langle \alpha, \beta \rangle$	$\mu > 0$
	$\sum_k \lambda_k f(\mu_k x)$	$(\sum_k \lambda_k \mu_k^{-s}) \cdot f^*(s)$		Th. 1, 5, Lem. 1
F_4	$f(x) \log x$	$\frac{d}{ds} f^*(s)$	$\langle \alpha, \beta \rangle$	
F_5	$\Theta f(x)$	$-s f^*(s)$	$\langle \alpha', \beta' \rangle$	$\Theta = x \frac{d}{dx}$
	$\frac{d}{dx} f(x)$	$-(s-1) f^*(s-1)$	$\langle \alpha' - 1, \beta' - 1 \rangle$	
	$\int_0^x f(t) dt$	$-\frac{1}{s} f^*(s+1)$		

FIGURE 1. Basic functional properties of Mellin transforms. The table lists the original function, its Mellin transform, and the validity strips.

A particular case of the rule for powers is

$$\mathcal{M}[f(\frac{1}{x}); s] = -f^*(-s).$$

Since the function $1/x$ exchanges 0 and ∞ , this permits to limit consideration of asymptotic properties of a function to one of the two places $0, +\infty$. A useful application of the rule is for the transform of the Gaussian function,

$$\mathcal{M}[e^{-x^2}; s] = \frac{1}{2} \Gamma(\frac{s}{2}),$$

which arises in estimates of sums involving binomial coefficients.

The formal rule

$$\frac{d}{ds} f^*(s) = \int_0^\infty f(x)(\log x) x^{s-1} dx$$

is readily justified analytically by “differentiation under the integral sign”. (As complex-differentiable implies analytic, this also supports our earlier claim that Mellin transforms are analytic in the fundamental strip.) Thus

$$\mathcal{M}[f(x)(\log x); s] = \frac{d}{ds} f^*(s).$$

For instance, the transform of $H(x) \log x$ is $-1/s^2$, the transform of $e^{-x} \log x$ is $\Gamma'(s)$.

Conversely, the rule for transforming the derivative of an original function is best enunciated in terms of the operator

$$\Theta = x \frac{d}{dx}.$$

Frequently the fundamental strips of $f(x)$ and $\Theta f(x)$ have a nonempty intersection $\langle \alpha', \beta' \rangle$. This is ensured in the common case where $f(x) = \mathcal{O}(x^{-\alpha})$ as $x \rightarrow 0$ and $f(x) = \mathcal{O}(x^{-\beta})$ as $x \rightarrow \infty$ whenever $\Theta f(x)$ satisfies the same estimates (i.e., the asymptotic form of the derivative of f coincides with the derivative of the asymptotic form of f). Under this sufficient condition, with $f(x)$ continuous and piecewise differentiable, integration by parts yields

$$\int_0^{+\infty} \Theta f(x) x^{s-1} dx = [f(x) x^s]_0^{+\infty} - s \int_0^{+\infty} f(x) x^{s-1} dx \quad \alpha < \Re(s) < \beta,$$

and the term $[f(x) x^s]_0^\infty$ equals 0 for $s \in \langle \alpha, \beta \rangle$. Thus,

$$\mathcal{M}[\Theta f(x); s] = -s f^*(s).$$

This permits to deduce Mellin transforms of primitive functions by identification. For instance, from the transform of $(1+x)^{-1}$, we find

$$\mathcal{M}[\log(1+x), s] = \frac{\pi}{s \sin \pi s}, \quad -1 < \Re(s) < 0.$$

In the same vein,

$$\mathcal{M}[e^{-x} - 1, s] = \frac{\Gamma(s+1)}{s} = \Gamma(s), \quad -1 < \Re(s) < 0,$$

and so on, see Fig. 2.

Note on transforms of analytic functions. Although we do not make use of it here, there is a fruitful approach to the determination of transforms of analytic functions that is based on a classical loop integral representation due to Hankel [83, 85]. A *Hankel contour* \mathcal{H} is a simple loop that starts in the upper half-plane near $+\infty$, circles around the origin counterclockwise and returns to $+\infty$ in the lower half-plane. If $f(x)$ is analytic in some open set that contains $[0, +\infty)$ and satisfies reasonable growth conditions, then *Hankel's formula* holds,

$$(14) \quad \mathcal{M}[f(x), s] = \frac{-1}{2i \sin \pi s} \int_{\mathcal{H}} f(w) (-w)^{s-1} dw, \quad 0 < \sigma < \beta.$$

The proof is based on shrinking the contour towards the real axis and using the fact that in the infinitesimal limit the integrand is $f(x) x^{s-1}$ multiplied by $(e^{-i\pi s} - e^{i\pi s}) = -2i \sin \pi s$. Hankel's formula permits to determine all transforms of rational functions as well as transforms of many meromorphic functions. See [77, 83, 85].

e^{-x}	$\Gamma(s)$	$\langle 0, +\infty \rangle$
$e^{-x} - 1$	$\Gamma(s)$	$\langle -1, 0 \rangle$
$e^{-x} - 1 + x$	$\Gamma(s)$	$\langle -2, -1 \rangle$
e^{-x^2}	$\frac{1}{2}\Gamma(\frac{1}{2}s)$	$\langle 0, +\infty \rangle$
$\frac{1}{(1+x)}$	$\frac{\pi}{\sin \pi s}$	$\langle 0, 1 \rangle$
$\log(1+x)$	$\frac{\pi}{s \sin \pi s}$	$\langle -1, 0 \rangle$
$H(x) \equiv 1_{0 < x < 1}$	$\frac{1}{s}$	$\langle 0, +\infty \rangle$
$x^\alpha (\log x)^k H(x)$	$\frac{(-1)^k k!}{(s + \alpha)^{k+1}}$	$\langle -\alpha, +\infty \rangle \quad k \in \mathbb{N}$

FIGURE 2. A microdictionary of Mellin transforms.

Inversion. With again $s = \sigma + it$ and the change of variables $x = e^{-y}$, the Mellin transform becomes a Fourier transform

$$f^*(s) = \int_0^\infty f(x)x^{s-1} dx = - \int_{-\infty}^{+\infty} f(e^{-y})e^{-\sigma y}e^{-ity} dy.$$

This links the two transforms, hence it provides directly inversion theorems for Mellin transforms. We cite here two main versions, one related to Lebesgue integration, the other to Riemann integration.

Theorem 2 (Inversion). (i) Let $f(x)$ be integrable with fundamental strip $\langle \alpha, \beta \rangle$. If c is such that $\alpha < c < \beta$ and $f^*(c + it)$ is integrable, then the equality

$$\frac{1}{2i\pi} \int_{c-i\infty}^{c+i\infty} f^*(s)x^{-s} ds = f(x)$$

holds almost everywhere. Moreover, if $f(x)$ is continuous, then equality holds everywhere on $(0, +\infty)$.

(ii) Let $f(x)$ be locally integrable with fundamental strip $\langle \alpha, \beta \rangle$ and be of bounded variation in a neighbourhood of x_0 . Then, for any c in the interval (α, β) ,

$$\lim_{T \rightarrow \infty} \frac{1}{2i\pi} \int_{c-iT}^{c+iT} f^*(s)x^{-s} ds = \frac{f(x_0^+) + f(x_0^-)}{2}.$$

Proof. See [18, 82, 86]. \square

Figure 2 presents a few classical original-transform pairs (f, f^*) . More can be found in standard tables of integral transforms [19, 66, 67] and in [61, 77].

2. THE FUNDAMENTAL CORRESPONDENCE

There is a very precise correspondence between the asymptotic expansion of a function at 0 (and ∞), and poles of its Mellin transform in a left (resp. right) half-plane. Each individual term in such an asymptotic expansion of $f(x)$ having the form $x^c(\log x)^k$ is associated to a pole of $f^*(s)$ at $s = -c$ with multiplicity $k+1$. The correspondence fares both ways. It is thus the basis of an asymptotic process: *For estimating asymptotically a function $F(x)$ —typically a harmonic sum—, determine its Mellin transform and translate back its singularities into asymptotic terms in the expansion of $F(x)$.*

Let $\phi(s)$ be meromorphic at $s = s_0$: it admits near s_0 a *Laurent expansion*

$$(15) \quad \phi(s) = \sum_{k \geq -r} c_k (s - s_0)^k.$$

The function $\phi(s)$ has a pole of order r if $r > 0$ and $c_{-r} \neq 0$, it is analytic (regular, holomorphic) at s_0 if $r = 0$. A *singular element* of $\phi(s)$ at s_0 is an initial sum of the Laurent expansion (15) truncated at terms of order $\mathcal{O}(1)$ or smaller.

Definition 2 (Singular expansion). Let $\phi(s)$ be meromorphic in Ω with S including all the poles of $\phi(s)$ in Ω . A *singular expansion* of $\phi(s)$ in Ω is a formal sum of singular elements of $\phi(s)$ at all points of S .

When E is a singular expansion of $\phi(s)$ in Ω , we write

$$\phi(s) \asymp E \quad (s \in \Omega).$$

For instance, one has

$$(16) \quad \frac{1}{s^2(s+1)} \asymp \left[\frac{1}{s+1} + 2 + 3(s+1) \right]_{s=-1} + \left[\frac{1}{s^2} - \frac{1}{s} \right]_{s=0} + \left[\frac{1}{2} \right]_{s=1} \quad (s \in \langle -2, +2 \rangle),$$

where the point of expansion may be indicated whenever needed as a subscript to the corresponding singular element. The expansion (16) is a concise way of combining information contained in the Laurent expansions of the function $\phi(s) \equiv s^{-2}(s+1)^{-1}$ at the three points of $S = \{-1, 0, 1\}$:

$$\phi(s)_{s=-1} \equiv (s+1)^{-1} + 2 + 3(s+1) + \dots, \quad \phi(s)_{s=0} \equiv s^{-2} - s^{-1} + 1 + \dots,$$

$$\phi(s)_{s=1} \equiv \frac{1}{2} - \frac{5}{4}(s-1) + \frac{17}{8}(s-1)^2 + \dots.$$

EXAMPLE 1. The Gamma function. The Mellin transform of the function e^{-x} defines the Gamma function,

$$\Gamma(s) = \int_0^\infty e^{-x} x^{s-1} dx,$$

for $\Re(s) > 0$. Integration by part permits to verify the well known functional equation

$$\Gamma(s+1) = s\Gamma(s),$$

which allows to extend $\Gamma(s)$ to a meromorphic function in the whole of \mathbb{C} . The function $\Gamma(s)$ satisfies $\Gamma(1) = 1$, and from the functional equation, one has

$$\Gamma(s) = \frac{\Gamma(s+m+1)}{s(s+1)(s+2)\cdots(s+m)}.$$

Thus, $\Gamma(s)$ has poles at all the points $s = -m$ with $m \in \mathbb{N}$, near which

$$\Gamma(s) \sim \frac{(-1)^m}{m!} \frac{1}{s+m},$$

so that the Gamma function admits

$$(17) \quad \Gamma(s) \asymp \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \frac{1}{s+k} \quad (s \in \mathbb{C})$$

as singular expansion in the whole of \mathbb{C} . □

Direct mapping. The function e^{-x} has a Taylor expansion at $x = 0$:

$$(18) \quad e^{-x} = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} x^k.$$

There is a striking coincidence of coefficients in the Taylor expansion (18) of the original function e^{-x} and in the singular expansion (17) of the transform $\Gamma(s)$ expressed by the rule

$$x^k \mapsto \frac{1}{s+k}.$$

This is in fact a completely general phenomenon.

Theorem 3 (Direct mapping). *Let $f(x)$ have a transform $f^*(s)$ with non-empty fundamental strip $\langle \alpha, \beta \rangle$.*

(i) *Assume that $f(x)$ admits as $x \rightarrow 0^+$ a finite asymptotic expansion of the form*

$$(19) \quad f(x) = \sum_{(\xi, k) \in A} c_{\xi, k} x^{\xi} (\log x)^k + \mathcal{O}(x^{\gamma})$$

where the ξ satisfy $-\gamma < -\xi \leq \alpha$ and the k are nonnegative. Then $f^(s)$ is continuable to a meromorphic function in the strip $\langle -\gamma, \beta \rangle$ where it admits the singular expansion*

$$f^*(s) \asymp \sum_{(\xi, k) \in A} c_{\xi, k} \frac{(-1)^k k!}{(s+\xi)^{k+1}} \quad (s \in \langle -\gamma, \beta \rangle).$$

(ii) *Similarly, assume that $f(x)$ admits as $x \rightarrow +\infty$ a finite asymptotic expansion of the form (19) where now $\beta \leq -\xi < -\gamma$. Then $f^*(s)$ is continuable to a meromorphic function in the strip $\langle \alpha, -\gamma \rangle$ where*

$$f^*(s) \asymp - \sum_{(\xi, k) \in A} c_{\xi, k} \frac{(-1)^k k!}{(s+\xi)^{k+1}} \quad (s \in \langle \alpha, -\gamma \rangle).$$

Thus terms in the asymptotic expansion of $f(x)$ at 0 induce poles of $f^*(s)$ in a strip to the *left* of the fundamental strip; terms in the expansion at $+\infty$ induce poles in a strip to the *right*.

Proof. (See [18].) Since $\mathcal{M}(f(1/x), s) = -\mathcal{M}(f(x), -s)$, it suffices to treat the case (i) corresponding to $x \rightarrow 0^+$. By assumption, the function $g(x)$

$$g(x) = f(x) - \sum_{(\xi, k) \in A} c_{\xi, k} x^{\xi} (\log x)^k$$

satisfies $g(x) = \mathcal{O}(x^{\gamma})$.

For s in the fundamental strip, a split of the definition domains yields

(20)

$$\begin{aligned} f^*(s) &= \int_0^1 f(x) x^{s-1} dx + \int_1^{\infty} f(x) x^{s-1} dx \\ &= \int_0^1 g(x) x^{s-1} dx + \int_0^1 \left(\sum_{(\xi, k) \in A} c_{\xi, k} x^{\xi} (\log x)^k \right) x^{s-1} dx + \int_1^{+\infty} f(x) x^{s-1} dx. \end{aligned}$$

In the last line of (20), the first integral defines an analytic function of s in the strip $\langle -\gamma, +\infty \rangle$ since $g(x) = \mathcal{O}(x^{\gamma})$ as $x \rightarrow 0$; the third integral is analytic in $\langle -\infty, \beta \rangle$, so that the sum of these two is analytic in $\langle -\gamma, \beta \rangle$. Finally, straight integration expresses the middle integral as

$$\sum_{(\xi, k) \in A} c_{\xi, k} \frac{(-1)^k k!}{(s + \xi)^{k+1}}$$

which is meromorphic in all \mathbb{C} and provides the singular expansion of $f^*(s)$ in the extended strip. \square

In the case where there exists a complete expansion of $f(x)$ at 0 (or ∞), the transform $f^*(s)$ becomes analytic in a complete left (or right) half-plane. This situation always occurs for functions that are analytic at 0 (or $+\infty$).

EXAMPLE 2. *The zeta function.* We gave earlier the Mellin pair

$$f(x) = \frac{e^{-x}}{1 - e^{-x}}, \quad f^*(s) = \Gamma(s)\zeta(s)$$

with fundamental strip $\langle 1, \infty \rangle$. The function $f(x)$ is exponentially small at infinity and it admits a complete expansion near $x = 0$

$$\frac{e^{-x}}{1 - e^{-x}} = \sum_{k=-1}^{\infty} B_{k+1} \frac{x^k}{(k+1)!},$$

which defines the Bernoulli numbers B_k : $B_0 = 1$, $B_1 = -\frac{1}{2}$, etc. Thus, $f^*(s) = \Gamma(s)\zeta(s)$ is meromorphic in the whole of \mathbb{C} with singular expansion

$$(21) \quad \Gamma(s)\zeta(s) \asymp \sum_{k=-1}^{\infty} \frac{B_{k+1}}{(k+1)!} \frac{1}{s+k}.$$

Therefore, $\zeta(s)$ is meromorphic in the whole of \mathbb{C} (this is often proved by means of its functional equation). In addition comparison of the singular expansions of $\Gamma(s)$ in (17) and $\Gamma(s)\zeta(s)$ in (21) yields

$$\zeta(s) \underset{s \rightarrow -1}{\sim} \frac{1}{s-1}, \quad \zeta(0) = -\frac{1}{2}, \quad \zeta(-m) = -\frac{B_{m+1}}{m+1}.$$

□

EXAMPLE 3. *The transform of $(1+x)^{-1}$.* The function $f(x) = (1+x)^{-1}$ has $\langle 0, 1 \rangle$ as its fundamental strip. The two expansions,

$$\frac{1}{1+x} = \sum_{n=0}^{\infty} (-1)^n x^n \quad (x \rightarrow 0^+) \quad \text{and} \quad \frac{1}{1+x} = \sum_{n=1}^{\infty} (-1)^{n-1} x^{-n} \quad (x \rightarrow +\infty),$$

translate into

$$f^*(s) \asymp \sum_{n=0}^{\infty} \frac{(-1)^n}{s+n} \quad (s \in \langle -\infty, 1 \rangle) \quad \text{and} \quad f^*(s) \asymp - \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{s-n} \quad (s \in \langle 0, +\infty \rangle),$$

which is consistent with the known form,

$$f^*(s) \equiv \frac{\pi}{\sin \pi s} \asymp \sum_{n \in \mathbb{Z}} \frac{(-1)^n}{s+n} \quad (s \in \mathbb{C}).$$

□

The next example illustrates the fact that Mellin analysis may be applied to functions without explicit transforms.

EXAMPLE 4. *A non-explicit transform.* The function $f(x) = (\cosh(x))^{-1/2}$ is exponentially small at ∞ and near 0 it satisfies

$$\frac{1}{\sqrt{\cosh(x)}} = 1 - \frac{1}{4}x^2 + \frac{7}{96}x^4 - \frac{139}{5760}x^6 + \frac{5473}{645120}x^8 + \mathcal{O}(x^{10}),$$

so that its transform $f^*(s)$ is meromorphic in \mathbb{C} and

$$f^*(s) \asymp \frac{1}{s} - \frac{1}{4} \frac{1}{s+2} + \frac{7}{96} \frac{1}{s+4} - \frac{139}{5760} \frac{1}{s+6} + \frac{5473}{645120} \frac{1}{s+8} + \dots$$

□

A general principle also derives from the proof of Theorem 3: *subtracting from a function a truncated form of its asymptotic expansion at either 0 or ∞ does not alter its Mellin transform and only shifts the fundamental strip.* An instance is provided by the equalities

$$(22) \quad \mathcal{M}(e^{-x}, s) = \Gamma(s) \quad s \in \langle 0, +\infty \rangle, \quad \mathcal{M}(e^{-x} - 1, s) = \Gamma(s) \quad s \in \langle -1, 0 \rangle,$$

previously established using integration by parts and specific properties of the exponential. The following proof of (22) demonstrates the general technique on this particular example. Take the function

$$F^*(s) = \frac{1}{s} + \int_0^1 (e^{-x} - 1)x^{s-1} dx + \int_1^\infty e^{-x} x^{s-1} dx.$$

$f(x)$	$f^*(s)$
Order at 0: $\mathcal{O}(x^a)$	Leftmost boundary of f.s. at $\Re(s) = -a$
Order at $+\infty$: $\mathcal{O}(x^b)$	Rightmost boundary of f.s. at $\Re(s) = -b$
Expansion till $\mathcal{O}(x^\gamma)$ at 0	Meromorphic continuation till $\Re(s) \geq -\gamma$
Expansion till $\mathcal{O}(x^\delta)$ at ∞	Meromorphic continuation till $\Re(s) \leq -\delta$
Term $x^a(\log x)^k$ at 0	Pole with singular element $\frac{(-1)^k k!}{(s+a)^{k+1}}$
Term $x^a(\log x)^k$ at ∞	Pole with singular element $-\frac{(-1)^k k!}{(s+a)^{k+1}}$

FIGURE 3. The fundamental correspondence: aspects of the direct mapping (Theorem 3).

$f^*(s)$	$f(x)$
Pole at ξ left of fund. strip right of fund. strip	Term in asymptotic expansion $\approx x^{-\xi}$ expansion at 0 expansion at $+\infty$
Multiple pole left: $\frac{1}{(s-\xi)^{k+1}}$ right: $\frac{1}{(s-\xi)^{k+1}}$	Logarithmic factor $\frac{(-1)^k}{k!} x^{-\xi} (\log x)^{k-1}$ at 0 $-\frac{(-1)^k}{k!} x^{-\xi} (\log x)^{k-1}$ at ∞
Pole with imaginary part: $\xi = \sigma + it$ Regularly spaced poles	Fluctuations: $x^{-\xi} = x^{-\sigma} e^{it \log x}$ Fourier series in $\log x$

FIGURE 4. The fundamental correspondence: aspects of the converse mapping (Theorem 4 and Corollary 1).




FIGURE 5. The contour \mathcal{C} used in the proof of Theorem 4. Dots represent poles of $f^*(s)$.

Consideration of both integrals shows that the function is meromorphic in $\langle -1, +\infty \rangle$. Its restriction to $\langle 0, +\infty \rangle$ is

$$\Gamma(s) = \int_0^\infty e^{-x} x^{s-1} dx,$$

and its restriction to $\langle -1, 0 \rangle$ is

$$\int_0^\infty (e^{-x} - 1) x^{s-1} dx.$$

This argument shows that the transforms of e^{-x} and of $e^{-x} - 1$ are elements of the same meromorphic function in different strips.

Converse mapping. Under a set of mild conditions, a converse to the Direct Mapping theorem also holds: The singularities of a Mellin transform which is small enough towards $\pm i\infty$ encode the asymptotic properties of the original function.

Theorem 4 (Converse mapping). *Let $f(x)$ be continuous in $]0, +\infty[$ with Mellin transform $f^*(s)$ having a non-empty fundamental strip $\langle \alpha, \beta \rangle$.*

(i) *Assume that $f^*(s)$ admits a meromorphic continuation to the strip $\langle \gamma, \beta \rangle$ for some $\gamma < \alpha$ with a finite number of poles there, and is analytic on $\Re(s) = \gamma$. Assume also that there exists a real number $\eta \in (\alpha, \beta)$ such that*

$$(23) \quad f^*(s) = \mathcal{O}(|s|^{-r}) \quad \text{with } r > 1,$$

when $|s| \rightarrow \infty$ in $\gamma \leq \Re(s) \leq \eta$. If $f^*(s)$ admits the singular expansion for $s \in \langle \gamma, \alpha \rangle$

$$(24) \quad f^*(s) \asymp \sum_{(\xi, k) \in A} d_{\xi, k} \frac{1}{(s - \xi)^k},$$

then an asymptotic expansion of $f(x)$ at 0 is

$$f(x) = \sum_{(\xi, k) \in A} d_{\xi, k} \left(\frac{(-1)^{k-1}}{(k-1)!} x^{-\xi} (\log x)^k \right) + \mathcal{O}(x^{-\gamma}).$$

(ii) Similarly assume that $f^*(s)$ admits a meromorphic continuation to $\langle \alpha, \gamma \rangle$ for some $\gamma > \beta$ and is analytic on $\Re(s) = \gamma$. Assume also that the growth condition (23) holds in $\langle \eta, \gamma \rangle$ for some $\eta \in (\alpha, \beta)$. If $f^*(s)$ admits the singular expansion (24) for $s \in \langle \eta, \gamma \rangle$, then an asymptotic expansion of $f(x)$ at ∞ is

$$f(x) = - \sum_{(\xi, k) \in A} d_{\xi, k} \left(\frac{(-1)^{k-1}}{(k-1)!} x^{-\xi} (\log x)^k \right) + \mathcal{O}(x^{-\gamma}).$$

Proof. The proof makes use of the inversion theorem and of a residue computation using large rectangular contours in the extended strip of $f^*(s)$, see Fig. 5. As before, it suffices to consider case (i) corresponding to continuation to the left.

Let \mathcal{S} be the set of poles in $\langle \gamma, \beta \rangle$. Consider the integral

$$J(T) = \frac{1}{2i\pi} \int_{\mathcal{C}} f^*(s) x^{-s} ds,$$

where $\mathcal{C} \equiv \mathcal{C}(T)$ denotes the rectangular contour defined by the segments

$$[\eta - iT, \eta + iT], \quad [\eta + iT, \gamma + iT], \quad [\gamma + iT, \gamma - iT], \quad [\gamma - iT, \eta - iT].$$

Assume that T is larger than $|\Im(s_0)|$ for all poles $s_0 \in \mathcal{S}$. By Cauchy's theorem, $J(T)$ is equal to the sum of residues, which is by a direct computation

$$R = \sum_{(\xi, k) \in A} d_{\xi, k} \operatorname{Res} \left(\frac{x^{-s}}{(s - \xi)^k} \right)_{s=\xi} = \sum_{(\xi, k) \in A} d_{\xi, k} \left(\frac{(-1)^{k-1}}{(k-1)!} x^{-\xi} (\log x)^k \right).$$

Let now T tend to $+\infty$. The integral along the two horizontal segments is $\mathcal{O}(T^{-r})$ and thus tends to 0 as $T \rightarrow \infty$. The integral along the vertical line $\Re(s) = \eta$ that lies inside the fundamental strip tends to the inverse Mellin integral which converges given the decrease assumption on f^* and equals $f(x)$ by the inversion theorem (since $f(x)$ is continuous). The integral along the vertical line $\Re(s) = \gamma$ is bounded by a quantity of the form

$$\frac{1}{2\pi} \int_{\gamma-i\infty}^{\gamma+i\infty} |f^*(s)| |x^{-s}| |ds| = \mathcal{O}(1) \int_0^\infty \frac{x^{-\gamma} dt}{(1+t)^r} = \mathcal{O}(x^{-\gamma}),$$

given the growth assumption on f^* .

Thus, in the limit, $J(\infty)$ equals $f(x)$ plus a remainder term that is $\mathcal{O}(x^{-\gamma})$ plus the sum of residues that is of the stated form in x and $\log x$. \square

In many cases, $f^*(s)$ is meromorphic in a complete left or right half plane and satisfies the conditions of Theorem 4; then a complete asymptotic expansion for $f^*(s)$ results. Such an expansion may be either convergent or divergent. If divergent, the expansion is by necessity only asymptotic. If convergent it may in some cases represent the function exactly, but this cannot be a general phenomenon decidable from the series alone² as $f(x)$ and $f(x) + \varpi(x)$ have the same asymptotic expansions at 0 and ∞ whenever $\varpi(x)$ is a "flat" function like $\varpi(x) = e^{-(x+1/x)}$.

EXAMPLE 5. *The transform of $(1+x)^{-\nu}$.* Take $\nu > 0$, and consider the function

$$\phi(s) = \frac{\Gamma(s)\Gamma(\nu-s)}{\Gamma(\nu)}$$

that is analytic in the strip $\langle 0, \nu \rangle$. The singular expansion to the left of $\Re(s) = 0$ is

$$\phi(s) \asymp \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \frac{\Gamma(\nu+k-1)}{\Gamma(\nu)} \frac{1}{(s+k)}.$$

The problem here is the behaviour of the original function

$$f(x) = \frac{1}{2i\pi} \int_{\nu/2-i\infty}^{\nu/2+i\infty} \phi(s)x^{-s} ds.$$

as $x \rightarrow 0$. The decrease of $\phi(s)$ along vertical lines results from the decrease of the Gamma function. Thus, the conditions of Theorem 5 are satisfied. In this way, one finds, for any integer M :

$$f(x) = \sum_{k=0}^M \frac{(-1)^k}{k!} \frac{\Gamma(\nu+k-1)}{\Gamma(\nu)} x^k + \mathcal{O}(x^{M+1/2}).$$

The series on the right hand side is a truncation of the Taylor series of $(1+x)^{-\nu}$, by virtue of the binomial theorem. We have thus proved that $f(x) = (1+x)^{-\nu} + \varpi(x)$ for a function $\varpi(x)$ that decreases at 0 faster than any power of x .

Sharper estimates show that the remainder integral tends uniformly to 0 when $0 < x < 1$, so that the representation is in fact exact. We have thus found indirectly the Mellin pair

$$f(x) = \frac{1}{(1+x)^\nu}, \quad f^*(s) = \frac{\Gamma(s)\Gamma(\nu-s)}{\Gamma(\nu)}.$$

□

EXAMPLE 6. *A classical divergent series.* The function

$$\phi(s) = \Gamma(1-s) \frac{\pi}{\sin \pi s}$$

is analytic in the strip $\langle 0, 1 \rangle$. In $\Re(s) < 1$, it admits the singular expansion

$$\phi(s) \asymp \sum_{n=0}^{\infty} (-1)^n n! \frac{1}{s+n},$$

²A proof of an exact representation requires that the remainder integrals along vertical lines tend to zero; this needs a fast enough and uniform decrease of $f^*(s)$ along vertical lines.

which encodes for the original function the asymptotic expansion

$$f(x) \sim \sum_{n=0}^{\infty} (-1)^n n! x^n.$$

In fact $\phi(s)$ is the Mellin transform of the "harmonic integral"

$$f(x) = \int_0^{\infty} \frac{e^{-t}}{1+xt} dt,$$

whose consideration goes back to Euler. \square

Periodicities. A pole of f^* at a point $\xi = \sigma + it$ with a non-zero imaginary part induces a term of the form

$$x^{-\xi} = x^{-\sigma} e^{-it \log x},$$

which contains an oscillating component that is periodic in $\log x$ with period $2\pi/t$. Mellin transforms are precisely useful for quantifying many nonelementary fluctuation phenomena. In this context, there sometimes occur a vertical line of regularly spaced poles (see Section 6 below). A direct extension of Theorem 4 allows for the presence of infinitely many poles in a finite strip.

Corollary 1. *The conclusions of Theorem 4 remain valid assuming only a weaker form of the growth condition (17) along a denumerable set of horizontal segments $|\Im(s)| = T_j$ where $T_j \rightarrow +\infty$.*

Proof. Use the proof of Theorem 4 but restrict T to belong to the discrete set T_j which, by the growth condition imposed, must avoid the poles of $f^*(s)$. \square

In particular *regularly spaced poles of $f^*(s)$ along a vertical line correspond to a Fourier series in $\log x$* . For instance, simple poles of $f^*(s)$ at the $\chi_k = \sigma + 2ik\pi/\log B$ will arise from a component of the form $(1 - B^{-s})^{-1}$ in $f^*(s)$. The corresponding residues r_k induce for $f(x)$ the infinite sum

$$\pm \sum_{k \in \mathbb{Z}} r_k x^{-\xi_k} = \pm x^{-\sigma} \sum_{k \in \mathbb{Z}} r_k \exp(-2ik\pi \log_B x),$$

where the sign is '+' for poles left of the fundamental strip (expansion at 0), '-' otherwise (expansion at ∞).

3. HARMONIC SUMS

This section builds upon the functional properties of the Mellin transform and the fundamental correspondence to develop a framework adapted to the analysis of harmonic sums.

Definition 3 (Harmonic sums). *A sum of the form*

$$(25) \quad G(x) = \sum_k \lambda_k g(\mu_k x)$$

is called a harmonic sum. The λ_k are the amplitudes, the μ_k are the frequencies, and $g(x)$ is called the base function.

The Dirichlet series of the harmonic sum is the sum

$$(26) \quad \Lambda(s) = \sum_k \lambda_k \mu_k^{-s}.$$

In this paper, the frequencies either decrease to zero or increase to ∞ . By possibly considering $G(1/x)$ and changing the base function accordingly, we may always reduce ourselves to the single case $\mu_k \rightarrow +\infty$, which we assume in this section, unless otherwise stated.

The Mellin transform of a harmonic sum factors, as already mentioned in the introduction, into a product of the Dirichlet series $\Lambda(s)$ and of the transform $g^*(s)$ of the base function. This is true unconditionally for finite harmonic sums. Elementary arguments provide a first extension to infinite sums.

Lemma 2. Assume that $g(x)$ is bounded over any interval $[a, b] \subset (0, +\infty)$, and that it satisfies $g(x) = \mathcal{O}(x^u)$ when $x \rightarrow 0$ and $g(x) = \mathcal{O}(x^v)$ when $x \rightarrow +\infty$, with $u > v$. Assume that the Dirichlet series $\Lambda(s)$ has a half-plane of absolute convergence $\Re(s) > \sigma_a$ that has a non-empty intersection Δ with the strip $\langle -u, -v \rangle$.

Then the harmonic sum $G(x)$ of (25) is defined for all x in $(0, +\infty)$. The transform $G^*(s)$ is well-defined in Δ where it factors as

$$G^*(s) = \Lambda(s) \cdot g^*(s).$$

We recall that the theory of generalized Dirichlet series [41] guarantees that a Dirichlet series like $\Lambda(s)$ has a half-plane of absolute convergence $\Re(s) > \sigma_a$ and a half-plane of simple convergence $\Re(s) > \sigma_c$ where

$$\sigma_a - \sigma_c \geq 0.$$

Proof. First consider the special case where $g(x)$ is $\mathcal{O}(1)$ at 0, is $\mathcal{O}(x^{-1})$ at ∞ , and $\Lambda(s)$ converges absolutely for $\Re(s) > -\delta$ for some $\delta > 0$.

Then over $(0, +\infty)$, we have both $|g(x)| < C/x$ and $|g(x)| < D$, for some constants $C, D > 0$. Thus, one has

$$|G(x)| < D \cdot \sum_k |\lambda_k| < \infty,$$

by the assumption that $\Lambda(0)$ converges absolutely. Consequently, $G(x) = \mathcal{O}(1)$ everywhere and in particular at 0. By summation, $G(x)$ is also $\mathcal{O}(x^{-1})$ everywhere, and in particular as $x \rightarrow +\infty$. By the dominated convergence theorem, $G^*(s)$ therefore exists in the strip $\langle 0, 1 \rangle$.

The general case reduces to the special case by normalizing $g(x)$ and considering

$$\hat{g}(x) = x^{-u/(v-u)} g(x^{1/(v-u)}).$$

□

Our treatment throughout this paper relies on the assumption that $g^*(s)$ and $\Lambda(s)$ in Lemma 2 are continuable as meromorphic functions in regions of the complex plane larger than what their definitions imply, and additionally satisfy controlled growth conditions towards $\pm i\infty$.

Definition 4. (i) A function $\phi(s)$ is said to be of fast decrease in the closed strip $\sigma_1 \leq \Re(s) \leq \sigma_2$ if for any $r > 0$,

$$\phi(s) = \mathcal{O}(|s|^{-r}),$$

as $|s| \rightarrow \infty$ in the strip.

(ii) A function is said to be of slow increase in the closed strip $\sigma_1 \leq \Re(s) \leq \sigma_2$ if for some $r > 0$,

$$\phi(s) = \mathcal{O}(|s|^r),$$

as $|s| \rightarrow \infty$ in the strip.

The property of fast decrease means that $\phi(s)$ decays faster than any negative power of $|s|$. For instance, the function $\pi/\sin \pi s$ is of fast decrease in any finite closed strip since the complex exponential representation yields

$$\frac{\pi}{\sin \pi s} = \mathcal{O}(e^{-\pi|t|}), \quad s = \sigma + it,$$

which exhibits an exponential decrease along vertical lines. A similar exponential decay holds for $\Gamma(s)$ by the complex version of Stirling's formula [1, p. 257]:

$$(27) \quad |\Gamma(\sigma + it)| \sim \sqrt{2\pi} |t|^{\sigma-1/2} e^{-\pi|t|/2} \quad (t \rightarrow +\infty).$$

The property of slow increase holds for many Dirichlet series and for $\sigma < 1$, one has [83, p. 94]

$$(28) \quad \zeta(\sigma + it) = \mathcal{O}(|t|^{1-\sigma}).$$

(Uniform versions of (27), (28) also exist, see the cited references.)

The property of slow increase is also called “finite order”. It means that $\phi(s)$ is of at most polynomial growth in the strip. The function $\zeta(s)$ is bounded in any closed strip of the form $[1 + \epsilon, M]$ and thus of slow increase there. Classical theorems [83] that extend (28) near $\Re(s) = 1$ show that it is of slow increase in any finite strip of the complex plane.

The theorem below represents the basic paradigm for the analysis of harmonic sums. It presupposes that *the Mellin transform of the base function is of fast decrease and the Dirichlet series of the harmonic sum is of moderate growth in an extended region of the complex plane*. In the statement, only the abscissa of simple convergence is needed, a handy improvement for many applications.

Theorem 5 (Harmonic sums). Let $g^*(s)$ have a fundamental strip $\langle \alpha, \beta \rangle$. Let $\Lambda(s)$ admit the half-plane of simple convergence $\Re(s) > \sigma_c$. Assume that

- the half-plane of convergence of $\Lambda(s)$ intersects the fundamental strip of $g^*(s)$: $\sigma_c < \beta$ and let $\alpha' = \max(\alpha, \sigma_c)$;
- the functions $g^*(s)$ and $\Lambda(s)$ admit a meromorphic continuation in a strip $\langle \gamma, \beta \rangle$ and are analytic on $\Re(s) = \gamma$, for some $\gamma < \alpha$;
- on the closed strip $\gamma \leq \Re(s) \leq (\alpha' + \beta)/2$, the function $g^*(s)$ is of fast decrease and the function $\Lambda(s)$ is of slow increase.

Then the harmonic sum $G(x)$ converges for all $x > 0$ on $(0, +\infty)$. An asymptotic expansion of $G(x)$ as $x \rightarrow 0$ till an error term $\mathcal{O}(x^{-\gamma})$ is obtained by termwise translation of the singular expansion of $G^*(s) = \Lambda(s)g^*(s)$ according to the rule

$$\frac{A}{(s-\xi)^{k+1}} \mapsto A \frac{(-1)^k}{k!} x^{-\xi} (\log x)^k.$$

Proof. First select an arbitrary integration abscissa σ in (α', β) and take some σ_0 such that $\alpha' < \sigma_0 < \sigma$. The inversion theorem provides

$$(29) \quad \sum_{n=1}^N \lambda_n g(\mu_n x) = \int_{\sigma-i\infty}^{\sigma+i\infty} \sum_{n=1}^N \frac{\lambda_n}{\mu_n^s} g^*(s) x^{-s} ds.$$

There is a well-known bound on the growth of a (generalized) Dirichlet series in any fixed half-plane strictly interior to its simple convergence domain that reads

$$|\Lambda(s)| \leq C(|s| + 1),$$

for some constant C , see [41, p. 3-4]. From this, we have

$$(30) \quad \left| \sum_{n=1}^N \frac{\lambda_n}{\mu_n^s} g^*(s) x^{-s} \right| \leq C(|s| + 1) \cdot |g^*(s)| \cdot x^{-\sigma},$$

which permits to apply the dominated convergence theorem and establishes the convergence of $G(x)$ expressed as an inverse Mellin integral:

$$(31) \quad G(x) = \frac{1}{2i\pi} \int_{\sigma-i\infty}^{\sigma+i\infty} \Lambda(s) g^*(s) x^{-s} ds.$$

In addition, the bound (30) shows that $G(x)$ is $\mathcal{O}(x^{-\sigma})$ where σ was chosen arbitrarily in (α', β) . Thus, the strip (α', β) is included in the fundamental strip of $G(x)$.

The upper bound of $\mathcal{O}(x^{-\sigma})$ also holds for all the partial sums $\sum_{n=1}^N \lambda_n g(\mu_n x)$ which are therefore dominated by an integrable function over $(0, +\infty)$. Thus, the dominated convergence theorem applies once more and yields

$$G^*(s) = \lim_{N \rightarrow \infty} \int_0^\infty \sum_{n=1}^N \lambda_n \cdot g(\mu_n x) x^{s-1} dx = \left(\lim_{N \rightarrow \infty} \sum_{n=1}^N \frac{\lambda_n}{\mu_n^s} \right) \cdot g^*(s),$$

that is to say the fundamental relation $G^*(s) = \Lambda(s)g^*(s)$.

The proof is now completed by a straight application of the *Converse Mapping Theorem* (Theorem 4). \square

A symmetric result holds near $x \rightarrow \infty$, assuming now the meromorphic extensions to hold to the left of the fundamental strip; the sign is reversed due to the orientation of the contour. This discussion is summarized by the following informal statement.

Mellin summation formula. Under the conditions of Theorem 5,

$$\sum_k \lambda_k g(\mu_k x) \sim \pm \sum_{s \in H} \text{Res}(g^*(s) \Lambda(s) x^{-s}),$$

- where, for an expansion near 0, the sum is over the set H of poles to the left of the fundamental strip, and the sign is '+';
- for an expansion near ∞ , the sum is over poles to the right of the fundamental strip, and the sign is '-'.

In addition, the relation (31) shows that $G(x)$ is of C^∞ class since $\Lambda(s)g^*(s)$ is of fast decrease so that derivation under the integral sign is permitted.

EXAMPLE 7. *Harmonic numbers.* The function

$$h(x) = \sum_{k=1}^{\infty} \left[\frac{1}{k} - \frac{1}{k+x} \right] = \sum_{k=1}^{\infty} \frac{1}{k} \frac{x}{1 + \frac{x}{k}},$$

satisfies $h(n) = H_n$ where H_n is the harmonic number. (It is closely related to $\frac{d}{dx} \log \Gamma(x)$.) Its Mellin transform is

$$h^*(s) = -\frac{\pi}{\sin \pi s} \zeta(1-s),$$

with fundamental strip $\langle -1, 0 \rangle$ and singular expansion to the right of this fundamental strip

$$h^*(s) \asymp \left[\frac{1}{s^2} - \frac{\gamma}{s} \right] - \sum_{k=1}^{\infty} (-1)^k \frac{\zeta(1-k)}{s-k}.$$

Thus (with a minus sign corresponding to the expansion at ∞)

$$H_n \sim \log n + \gamma + \frac{1}{2n} + \sum_{k \geq 2} \frac{(-1)^k B_k}{k} \frac{1}{n^k} \sim \log n + \gamma + \frac{1}{2n} - \frac{1}{12n^2} + \frac{1}{120n^4} + \dots$$

Generalized harmonic numbers can be analyzed in the same way, see [37, p. 300] for a table giving $H_n^{(2)}$. \square

EXAMPLE 8. *Stirling's formula.* From the product decomposition of the Gamma function, one has

$$\ell(x) := \log \Gamma(x+1) - \gamma x = \sum_{n=1}^{\infty} \left[\frac{x}{n} - \log \left(1 + \frac{x}{n} \right) \right], \quad (s \in \langle -2, +\infty \rangle).$$

This example is similar in spirit to the previous one. The Mellin transform is

$$\ell^*(s) = -\zeta(-s) \frac{\pi}{s \sin \pi s},$$

with fundamental strip $\langle -2, -1 \rangle$. There are double poles at $s = -1$, $s = 0$ and simple poles at the positive integers,

$$\ell^*(s) \asymp \left[\frac{1}{(s+1)^2} + \frac{1-\gamma}{(s+1)} \right] + \left[\frac{1}{2s^2} - \frac{\log \sqrt{2\pi}}{s} \right] + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} \zeta(-n)}{n(s+n)}.$$

Hence Stirling's formula

$$\log(x!) \sim \log \left(x^x e^{-x} \sqrt{2\pi x} \right) + \sum_{n=1}^{\infty} \frac{B_{2n}}{2n(2n-1)} \frac{1}{x^{2n-1}}.$$

\square

EXAMPLE 9. *Polylogarithms.* The function

$$L_w(x) = \sum_{n=1}^{\infty} \frac{e^{-nx}}{n^w},$$

is related to polylogarithms that are defined for integer w by $\text{Li}_w(z) = \sum_{n=1}^{\infty} z^n n^{-w}$. For $w \in \mathbb{N}$, one has $L_w(x) = \text{Li}_w(e^{-x})$. We assume first that w is a positive integer, $w = k$.

The function $L_k(x)$ is a typical harmonic sum with transform

$$L_k^*(s) = \zeta(s+k)\Gamma(s),$$

valid in $\Re(s) > 1$ at least. The singular expansion obtains from the singular expansions of $\zeta(s)$ (shifted) and $\Gamma(s)$:

$$L_k^*(s) \asymp \sum_{\substack{n=0 \\ n \neq k-1}}^{\infty} (-1)^n \frac{\zeta(k-n)}{n!} \frac{1}{s+n} + \frac{(-1)^{k-1}}{(k-1)!} \left[\frac{1}{(s+k-1)^2} + \frac{H_{k-1}}{s+k-1} \right], \quad (s \in (-\infty, +\infty)).$$

The double pole at $s = k-1$ induces a logarithmic factor and

$$L_k(x) = \frac{(-1)^{k-1}}{(k-1)!} x^{k-1} [-\log x + H_{k-1}] + \sum_{\substack{n=0 \\ n \neq k-1}}^{\infty} (-1)^n \frac{\zeta(k-n)}{n!} x^n.$$

The right hand side is *a priori* only an asymptotic representation valid as $x \rightarrow 0$, but here the representation is exact as seen from uniform bounds deriving from growth formulæ for the Gamma and zeta functions (27,28). We have obtained in this way the Cohen-Zagier representation formula for polylogarithms [58, p. 387].

When w is not an integer, there is no confluence of singularities and accordingly, the logarithmic term disappears. For instance:

$$L_{1/2}(x) \equiv \sum_{n=1}^{\infty} \frac{e^{-nx}}{\sqrt{n}} = \sqrt{\frac{\pi}{x}} + \sum_{n=0}^{\infty} (-1)^n \frac{\zeta(\frac{1}{2}-n)}{n!} x^n.$$

□

EXAMPLE 10. *Modified theta functions.* The sum

$$\Theta_w(x) = \sum_{n=1}^{\infty} \frac{e^{-n^2 x^2}}{n^w}$$

is related to theta functions. The general case can be treated like before but we restrict attention to $w = 0, 1$. The case $w = 1$ arises in the analysis of bubble sort, see [56] p. 129 and Ex. 5.2.2-51. One has

$$\Theta_1^*(s) = \frac{1}{2} \Gamma\left(\frac{s}{2}\right) \zeta(s+1).$$

There is a double pole at $s = 0$ and poles of $\Gamma(s/2)$ are the even negative integers. This gives the expansion as $x \rightarrow 0$:

$$\begin{aligned} \Theta_1^*(s) &\asymp \left[\frac{1}{s^2} + \frac{\gamma}{2s} \right] + \frac{1}{12} \frac{1}{s+2} + \frac{1}{240} \frac{1}{s+4} + \dots \\ \Theta_1(x) &= -\log x + \frac{\gamma}{2} + \frac{1}{12} x^2 + \frac{1}{240} x^4 + \dots \end{aligned}$$

The situation for $\Theta_0(x)$ is different since the poles of $\Gamma(s/2)$ at the even integers are cancelled by the values of the zeta function: $\zeta(-2) = \zeta(-4) = \dots = 0$. Thus

$$\Theta_0(x) = \frac{1}{2} \frac{\sqrt{\pi}}{x} - \frac{1}{2} + \mathcal{O}(x^M), \quad (x \rightarrow 0)$$

for any $M > 0$. The degenerate asymptotic series can only be asymptotic. \square

This last example is related to the modular transformation of theta functions (see also Poisson's summation formula in Section 4), and various related forms have been investigated from the asymptotic standpoint by Ramanujan, Berndt, De Bruijn, Knuth, and Gonnet, see [8], [15, p. 43], [36], [56, p. 129]. Ramanujan discusses some criteria that entail the existence of finite expansions [8, Chap. XV] for similar functions.

In general, the absence of singularities in a left half plane indicates a terminating expansion with exponentially small error term. (In many practical situations estimates of the error term result from applying the saddle point method to remainder integrals.)

The two examples that follow are motivated by the two types of harmonic sums mentioned in the introduction. One involves strongly fluctuating amplitudes resulting though in a smooth global behaviour. The other has geometrically increasing frequencies so that the Dirichlet series has a line of poles resulting, by Corollary 1, in an oscillating asymptotic behaviour expressed by a Fourier series.

EXAMPLE 11. *A divisor sum.* The analysis of the height of trees (see the introduction) suggests considering the sum

$$D(x) = \sum_{k=1}^{\infty} d(k) e^{-kx},$$

with $d(k)$ being the number of divisors of k . Given the equality

$$\sum_{k=1}^{\infty} \frac{d(k)}{k^s} = \zeta^2(s),$$

the Mellin transform of $D(x)$ is

$$D^*(s) = \Gamma(s) \zeta^2(s),$$

with fundamental strip $(1, +\infty)$. There are singularities at $s = 1$, $s = 0$, then at all the odd negative integers, so that

$$D^*(s) \asymp \left[\frac{1}{(s-1)^2} + \frac{\gamma}{s-1} \right] + \left[\frac{1}{4s} \right]_{s=0} - \sum_{k=0}^{\infty} \frac{(\zeta(-2k-1))^2}{(2k+1)!} \frac{1}{s+2k+1}.$$

This translates into the asymptotic expansion of $D(x)$ as $x \rightarrow 0$:

$$D(x) \sim \frac{1}{x} (-\log x + \gamma) + \frac{1}{4} - \sum_{k=0}^{\infty} \frac{(\zeta(-2k-1))^2}{(2k+1)!} x^{2k+1}.$$

Clearly sums of divisors can be treated in a similar way as their Dirichlet series are expressible in terms of the Riemann zeta function. Ramanujan discovered a number of related formulæ later treated by Berndt and Evans using Mellin transforms [8]. \square

From the proof of Theorem 5 what essentially matters is the *balance* between the growth of $\Lambda(s)$ and the decrease of $g^*(s)$. The asymptotic expansion remains valid as long as $G^*(s)$ is globally $\mathcal{O}(s^{-r})$ for some $r > 1$ in the extended strip (the C^∞ character mentioned above may get lost, though). Also, in accordance with Corollary 1, it is sufficient that the decrease of $G^*(s)$ hold along a discrete set of horizontal segments. This permits to cope with situations involving infinitely many imaginary poles corresponding to an oscillatory behaviour.

EXAMPLE 12. *A doubly exponential sum and periodicities.* The prototype of harmonic sums with a fluctuating behaviour is the function

$$G(x) = \sum_{k=0}^{\infty} e^{-x2^k},$$

whose behaviour is sought as $x \rightarrow 0$. The Mellin transform is

$$G^*(s) = \Gamma(s) \cdot \left(\sum_{k=0}^{\infty} (2^k)^{-s} \right) = \frac{\Gamma(s)}{1-2^{-s}}.$$

There are infinitely many poles of $\Lambda(s)$ at the points

$$\chi_k = \frac{2ik\pi}{\log 2}.$$

The Gamma function decreases fast as $\Im(s) \rightarrow \pm\infty$. The Dirichlet series $\Lambda(s) = (1-2^{-s})^{-1}$ stays bounded along horizontal segments that pass in-between poles at $|\Im(s)| = (2k+1)\pi/\log 2$, for k an integer. Thus, Corollary 1 to Theorem 4 applies here.

The residue of $\Lambda(s)$ is $1/(\log 2)$ by virtue of periodicity. Thus,

$$(32) \quad \frac{1}{1-2^{-s}} \asymp \sum_{k \in \mathbb{Z}} \frac{1}{\log 2} \frac{1}{s - \chi_k}, \quad (s \in \mathbb{C}).$$

Globally, the poles of $G^*(s)$ are thus a double pole at 0, single poles induced by $\Lambda(s)$ at the χ_k , and poles at the negative integers. Translating to $G(x)$ yields

$$G(x) = -\log_2(x) - \frac{\gamma}{\log 2} + \frac{1}{2} + Q(\log_2 x) + \sum_{n=1}^{\infty} \frac{1}{1-2^n} \frac{(-x)^n}{n!},$$

again an exact representation. There $Q(\log_2 x)$ condenses the contribution from the non-zero imaginary poles that reflects (32):

$$Q(\log_2 x) = \frac{1}{\log 2} \sum_{k \in \mathbb{Z} \setminus \{0\}} \Gamma(\chi_k) x^{-\chi_k} = \frac{1}{\log 2} \sum_{k \in \mathbb{Z} \setminus \{0\}} \Gamma(\chi_k) e^{-2ik\pi \log_2 x}.$$

Thus, after the logarithmic term, $G(x)$ contains a fluctuating term of order $\mathcal{O}(1)$ that is expressed as a Fourier series in $\log_2 x$ with explicit coefficients of a Gamma type.

An entirely similar method permits to extract fluctuations of a sum like

$$K(x) = \sum_{k=0}^{\infty} \left(1 - e^{-x/2^k}\right),$$

when $x \rightarrow +\infty$, see Prop. 2 below. \square

We observe that $|\Gamma(\pm 2i\pi/\log 2)| \doteq 0.54521 \cdot 10^{-6}$, and due to the fast decrease of $\Gamma(s)$ along imaginary lines, the fluctuating function $Q(\log_2 x)$ stays bounded by 10^{-6} . Such minute fluctuations constitute a common feature of many Mellin analyses.

A closely related example is analysed by Hardy (his function $\phi_2(x)$ in [40, p. 37]) who notes that Ramanujan failed to obtain an elementary proof of the prime number theorem because his argument neglected these tiny “wobbles” arising from imaginary poles.

4. SUMMATORY FORMULÆ

What we call here an *asymptotic* summatory formula is a formula that gives the asymptotic form of a class of harmonic sums when either the amplitude-frequency pair stays fixed (and the base function is taken in a wide class of functions) or the base function stays fixed (and the amplitude-frequency pair varies). The generality afforded by Theorem 5 permits to deduce a large number of summatory formulæ, a typical case being generalizations of the Euler-Maclaurin formula.

Euler-Maclaurin summation. The classical Euler-Maclaurin formula provides a full asymptotic expansion expressing the convergence of a Riemann sum to the corresponding integral. The expansion assumes the base function to be C^∞ and then involves the Bernoulli numbers. The formula has been extended by Barnes [6] and later revisited by Gonnet [36] in the case of certain functions singular at the origin.

Proposition 1 (Generalized Euler-Maclaurin summation). *Let $g(x)$ be such that*

$$g(x) \underset{x \rightarrow 0}{\sim} \sum_{k=0}^{\infty} c_k x^{\alpha_k},$$

for some increasing sequence $\{\alpha_k\}$ with $\alpha_0 > -1$. Assume that $g^(s)$ exists in $(-\alpha_0, \beta)$ for some $\beta > 1$ and is meromorphically continuable to a function of fast decrease in any finite strip of $(-\infty, \beta)$. Then*

$$G(x) \equiv \sum_{k=1}^{\infty} g(kx) \underset{x \rightarrow 0}{\sim} \frac{1}{x} \int_0^{\infty} g(x) dx + \sum_{k=0}^{\infty} c_k \zeta(-\alpha_k) x^{\alpha_k}.$$

Proof. The transform $G^*(s)$ is $g^*(s)\zeta(s)$ with residue at $s = 1$ that equals $g^*(1)$, itself the integral of $g(x)$. By Theorem 3, there are also poles of $g^*(s)$ and $G^*(s)$ at each $-\alpha_k$. The result then follows by Theorem 5 after taking into account the slow increase of $\zeta(s)$ in finite strips. \square

EXAMPLE 13. *A sum with half-integer exponents.* The sum

$$S_n = \sum_{k=1}^{\infty} \frac{1}{k^{1/2}(n+k)^{3/2}}$$

is an Euler-Maclaurin sum since

$$S_n = \frac{1}{n^2} G\left(\frac{1}{n}\right) \quad \text{where} \quad G(x) = \sum_{k=1}^{\infty} g(kx), \quad g(x) = \frac{1}{x^{1/2}(1+x)^{3/2}}.$$

The expansion of $g(x)$ near $x = 0$ involves half-integer exponents:

$$g(x) \sim \sum_{k=0}^{\infty} c_k x^{k-1/2}, \quad c_k = (-1)^k (k+1) 2^{-2k} \binom{2k+1}{k}.$$

In addition, $g^*(s)$ has fundamental strip $\langle 1/2, 5/2 \rangle$, and an explicit transform (deriving from the Eulerian Beta integral or from an earlier example)

$$g^*(s) = B\left(s - \frac{1}{2}, 2 - s\right) = \frac{\Gamma(s - 1/2)\Gamma(2 - s)}{\Gamma(3/2)},$$

which guarantees its fast decrease. Thus, a simple computation yields

$$S_n \sim \frac{2}{n} + \sum_{k=0}^{\infty} c_k \zeta\left(-k + \frac{1}{2}\right) \frac{1}{n^{k+3/2}}.$$

□

Related examples like

$$\sum_{k=1}^{n-1} \frac{1}{\sqrt{k(n-k)}},$$

motivated by the analysis of interpolation sequential search, are discussed in [36] and tabulated in [37, p. 298]. (The analysis is however more delicate, with $g^*(s)$ only decreasing like $s^{-1/2}$.)

EXAMPLE 14. *An exponential sum with square roots.* The function

$$G(x) = \sum_{n=1}^{\infty} e^{-\sqrt{n}x}$$

admits the expansion

$$G(x) \underset{x \rightarrow 0}{\sim} \frac{2}{x^2} - \frac{1}{2} + \sum_{k=1}^{\infty} \frac{(-1)^k}{k!} \zeta\left(-\frac{k}{2}\right) x^k,$$

since its Mellin transform is $\Gamma(s)\zeta(s/2)$.

□

Many similar summatory formulæ can be given under the assumptions of Prop. 1, like

$$\begin{aligned} \sum_{k=1}^{\infty} (-1)^{k-1} g(kx) &\sim \sum_{k=0}^{\infty} c_k (1 - 2^{1+\alpha_k}) \zeta(-\alpha_k) x^{\alpha_k} \\ \sum_{k=1}^{\infty} (\log k) g(kx) &\sim \frac{1}{x} \log \frac{1}{x} \int_0^{\infty} g(x) dx + \frac{1}{x} \int_0^{\infty} g(x) (\log x) dx \\ &\quad + \sum_{k=0}^{\infty} c_k \zeta'(-\alpha_k) x^{\alpha_k}. \end{aligned}$$

For instance, Ramanujan gives [8]

$$\sum_{k \geq 1} e^{-kx} \log k = \frac{1}{x} \left(\log \frac{1}{x} - \gamma \right) + \log \sqrt{2\pi} + \mathcal{O}(x).$$

Several examples of Ramanujan have been treated systematically by Berndt and Evans using Mellin transforms [9]. Functions with logarithmic singularities at 0 can also be dealt with in this way, see the papers by Barnes and Gonnet for alternative approaches [6, 36].

Note on formal Mellin analyses. In this range of problems, it is of interest to observe that asymptotic series follow *formally* from an exchange of summations in the harmonic sum, with the Dirichlet series coming out as a purely divergent series (!). This observation often constitutes a useful heuristic in the discovery of possible summation formulæ then rigorously established by Mellin transforms. An illustration is provided by the formula

$$G(x) \equiv \sum_{k=1}^{\infty} (-1)^k g((2k+1)x) \sim \sum_{k=0}^{\infty} c_k \eta(-\alpha_k) x^{\alpha_k}, \quad \text{where} \quad \eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^k}{(2k+1)^s}$$

is an entire “*L*-function” associated with a character modulo 4, see [12]. In effect, one has formally

$$\begin{aligned} G(x) &\stackrel{!}{=} \sum_m (-1)^m \left(\sum_k c_k x^{\alpha_k} (2m+1)^{\alpha_k} \right) \\ &\stackrel{!}{=} \sum_k c_k x^{\alpha_k} \left(\sum_{m=0}^{\infty} (-1)^k (2m+1)^{\alpha_k} \right) \\ &\stackrel{!}{=} \sum_{k=0}^{\infty} c_k \eta(-\alpha_k) x^{\alpha_k}. \end{aligned}$$

The Mellin transform justifies such formal manipulations while “explaining” the occurrence of additional terms like the integral in the Euler-Maclaurin summation.

Dyadic sums. Sums involving powers of 2, of the type

$$G_w(x) = \sum_{k \geq 0} 2^{-kw} g\left(\frac{x}{2^k}\right),$$

are particularly frequent in the analysis of algorithms and we call them *dyadic* sums. In applications x usually represents a large parameter.

Proposition 2 (Dyadic sums). *Let $g(x)$ be such that*

$$g(x) \underset{x \rightarrow \infty}{\sim} \sum_{k=0}^{\infty} d_k x^{-\beta_k},$$

for some increasing sequence $\{\beta_k\}$ with $\beta_0 > 0$. Assume that $g^(s)$ exists in $\langle \alpha, \beta_0 \rangle$ for some $\alpha < 0$ and is meromorphically continuable to a function of fast decrease in any finite strip of $\langle \alpha, +\infty \rangle$. Then*

$$G_0(x) \equiv \sum_{k \geq 0} g\left(\frac{x}{2^k}\right) \underset{x \rightarrow +\infty}{\sim} \frac{1}{\log 2} \sum_{k \in \mathbb{Z}} g^*\left(\frac{2ik\pi}{\log 2}\right) e^{-2ik\pi \log_2 x} + \sum_{k=0}^{\infty} \frac{d_k}{1-2^{\beta_k}} x^{-\beta_k}.$$

Proof. The transform of $G_0(x)$ is

$$G_0^*(s) = \frac{g^*(s)}{1-2^s}.$$

and it is readily subjected to Theorem 5 and Corollary 1. \square

Proposition 2 is characteristic of a large number of similar summation formulæ. Confluence of singularities will in general induce logarithmic factors.

EXAMPLE 15. *The standard dyadic sum.* Consider the analysis of $G_0(x)$ when now $g(\infty) \neq 0$ and

$$g(x) \underset{x \rightarrow +\infty}{\sim} \sum_{k=0}^{\infty} d_k x^{-k}.$$

(Thus $d_0 = g(\infty) \neq 0$.) There is now a double pole of $G_0^*(s)$ at $s = 0$ so that a two term expansion is required for $g^*(s)$ there. By the fundamental splitting, one has

$$g^*(s) = \frac{g(\infty)}{s} + \int_0^1 g(x) x^{s-1} dx + \int_1^{\infty} (g(x) - g(\infty)) x^{s-1} dx,$$

so that

$$g^*(s) = \frac{g(\infty)}{s} + \gamma[g] + \mathcal{O}(s) \quad \text{where} \quad \gamma[g] = \int_0^1 g(x) \frac{dx}{x} + \int_1^{\infty} (g(x) - g(\infty)) \frac{dx}{x}.$$

The constant $\gamma[g]$ is called the *Euler constant* of g since $\gamma[e^{-x} - 1] = -\gamma$. Thus,

$$G(x) \sim g(\infty) \log_2 x + \frac{1}{2} g(\infty) + \frac{\gamma[g]}{\log 2} + P(\log_2 x) + \sum_{k=1}^{\infty} \frac{d_k}{1-2^k} x^{-k},$$

for an explicitly determined periodic function $P(\cdot)$.

This example illustrates in passing the general technique for determining terms of the series expansion of a Mellin transform outside of its convergence strip by adapting the technique of subtracted singularities of Theorem 3. \square

Poisson's summation formula. There is a fruitful connexion between the classical Poisson summation formula [43, vol. 2, p. 271]

$$(33) \quad \sum_{k=-\infty}^{+\infty} f(kx) = \frac{1}{x} \sum_{k=-\infty}^{+\infty} \hat{f}\left(\frac{2\pi}{x}k\right), \quad \hat{f}(u) = \int_{-\infty}^{+\infty} f(t)e^{-2i\pi tu} du,$$

the functional equation of the Riemann zeta function, and other formulæ of analysis like the modular transformation of theta functions (this was first observed by Riemann in his original memoir) or the partial fraction expansion of the cotangent. That connexion can be based on the Mellin transform of well-chosen harmonic sums. We only provide here brief indications and refer for details to [55] and works cited therein.

The hyperbolic cotangent admits two equivalent expressions, one as a (harmonic) sum of exponentials, the other as a (harmonic sum) partial fraction decomposition:

$$(34) \quad \begin{aligned} \pi \coth(\pi x) &= \frac{1 + e^{-2\pi x}}{1 - e^{-2\pi x}} = 1 + 2 \sum_{n \geq 1} e^{-2n\pi x} \\ &= \frac{1}{x} + \sum_{n=1}^{\infty} \frac{x}{x^2 + n^2}. \end{aligned}$$

The functional equation of the Riemann zeta function,

$$(35) \quad \zeta(s) = 2^s \pi^{s-1} \sin \frac{\pi s}{2} \Gamma(1-s) \zeta(1-s),$$

appears to be the “image” under the Mellin transform of the identity between the two forms of (34), see [83, p. 24].

The proof of Poisson's summation formula then results from taking the Mellin transform of both sides of (33) and inserting the functional equation of $\zeta(s)$. This approach has the merit of establishing a general equivalence between summation formulæ of the Poisson type and functional equations of Dirichlet series [7, 42, 55]. Davies' book contains a direct illustration [13] with a proof of the modular transformation of the Dedekind η function.

Perron's formula. A collection of summatory formulæ, usually referred to as Perron's formula express partial sums of coefficients of a Dirichlet series as complex integrals of the inverse Mellin type applied to the Dirichlet series itself [41]. In our perspective, Perron's formula results from taking the step function $H(x)$ of (13) as the base function. The transform of $H(x)$ being $H^*(s) = 1/s$, one has formally for a Dirichlet series $\Lambda(s) = \sum_k \lambda_k \mu_k^{-s}$:

$$\sum_{\mu_k x < 1} \lambda_k \equiv \sum_k \lambda_k H(\mu_k x) = \frac{1}{2i\pi} \int_{c-i\infty}^{c+i\infty} \Lambda(s) x^{-s} \frac{ds}{s}.$$

Perron's formula is essential in analytic number theory, for instance in the proof of the prime number theorem. It has been recently employed for the analysis of cost functions of “divide-and-conquer” algorithms. There it permits to capture periodic fluctuations that are often of a fractal nature. For applications to mergesort and

maxima finding in computational geometry, see [24, 25]; for digital sums that occur in the analysis of several algorithms (sorting networks, register allocation), we refer to [26]. A related class of divide-and-conquer recurrences (with maximum) has been treated via Tauberian methods by Fredman and Knuth [35], and elementarily by Pippenger [69]. Note that the use of Perron's formula is often delicate since the transform $H^*(s)$ does not have good decay properties towards $\pm i\infty$.

II. Combinatorial Applications

In this part, we survey some of the major applications of Mellin asymptotics to combinatorial problems. Sums related to Catalan numbers form the subject of Section 5 and Section 6 develops the important "Bernoulli splitting process" that leads to dyadic sums. Additional combinatorial applications form the subject of Section 7.

5. CATALAN SUMS

Catalan sums have the form

$$(36) \quad S_n = \sum_{k=1}^n \lambda_k \frac{\binom{2n}{n-k}}{\binom{2n}{n}},$$

where the λ_k are of an arithmetical character. Such binomial sums occur in average values of characteristic parameters of combinatorial objects enumerated by the Catalan numbers,

$$(37) \quad C_n = \frac{1}{n+1} \binom{2n}{n},$$

like plane trees, binary trees, or ballot sequences [11, 38].

The historic paper of De Bruijn, Knuth, and Rice [16] provided the first analysis of this type. It concerns the expected height \overline{H}_n of rooted plane trees of n nodes under the uniform distribution. The sequence λ_k is then the divisor function $d(k)$. We briefly explain here the connection between this combinatorial problem and a Catalan sum like (36). (See also [51, p. 135] for details.)

A plane tree decomposes recursively as a root node to which is attached a sequence of trees. Let A_n be the number of trees with n nodes; the ordinary generating function of the sequence $\{A_n\}$ is defined by

$$A(z) = \sum_{n=1}^{\infty} A_n z^n.$$

Then, by the classical laws of combinatorial analysis (see, e.g., [33, 38, 78, 87]), the decomposition translates into a functional equation that admits an explicit solution

$$A(z) = \frac{z}{1 - A(z)} \quad \text{and} \quad A(z) = \frac{1 - \sqrt{1 - 4z}}{2}.$$

We have $A_{n+1} = C_n$ with C_n the Catalan number of (37).

Let similarly $A_h(z)$ be the generating function of trees of height at most h . As height is inherited from subtrees, one has then the basic recurrence

$$A_{h+1}(z) = \frac{z}{1 - A_h(z)} \quad \text{with} \quad A_1(z) = z.$$

Thus, the $A_h(z)$ are rational fractions that are also approximants to an infinite continued fraction representing $A(z)$. Solving the implied recurrence yields the closed form

$$A_h(z) = 2z \frac{(1 + \sqrt{1 - 4z})^h - (1 - \sqrt{1 - 4z})^h}{(1 + \sqrt{1 - 4z})^{h+1} - (1 - \sqrt{1 - 4z})^{h+1}}.$$

There results that the A_h can be expressed in terms of $A(z)$ alone; their Taylor expansion then derives by the Lagrange-Bürmann inversion theorem for analytic functions:

$$(38) \quad A_{n+1,h} - A_{n+1,h-1} = \sum_{j \geq 1} \omega_{jh}(n),$$

where

$$\omega_m(n) = \Delta^2 \binom{2n}{n-m} = \binom{2n}{n+1-m} - 2 \binom{2n}{n-m} + \binom{2n}{n-1-m}.$$

Thus, the number of trees of height $\geq h$ appears as a "sampled" sum the $2n$ th line of Pascal's triangle (upon taking second order differences).

By a well-known form of the expectations of discrete random variables, the mean height \overline{H}_{n+1} satisfies

$$\overline{H}_{n+1} = \frac{1}{A_{n+1}} \sum_h \sum_j \omega_{jh}(n).$$

Grouping terms according to the value of jh then reduces this expression to a simple sum:

$$(39) \quad \overline{H}_{n+1} = \frac{1}{A_{n+1}} \sum_k d(k) \omega_k(n).$$

We are therefore lead to considering sums of a pattern similar to (36),

$$S_n^{(a)} = \sum_{k=1}^n d(k) \frac{\binom{2n}{n-k-a}}{\binom{2n}{n}},$$

since

$$(40) \quad \frac{1}{n+1} \overline{H}_{n+1} = S_n^{(1)} - 2S_n^{(0)} + S_n^{(-1)},$$

the treatment of the central sum being typical. Stirling's formula yields the Gaussian approximation of binomial numbers: for $k = w\sqrt{n}$, and with $k = o(n^{3/4})$, one has

$$\frac{\binom{2n}{n-k}}{\binom{2n}{n}} \sim e^{-w^2} \left(1 - \frac{w^4 + 3w^2}{6n} + \frac{5w^8 + 6w^6 - 45w^4 - 60}{360n^2} + \dots \right).$$

This leads to introducing the continuous harmonic sum

$$G(x) = \sum_k d(k) e^{-k^2 x^2},$$

and an elementary argument (domination of the central terms) justifies the use of the Gaussian approximation inside $S_n^{(0)}$:

$$(41) \quad S_n^{(0)} = G\left(\frac{1}{\sqrt{n}}\right) + o(1).$$

The asymptotic analysis of $G(x)$ for small x (now, $x = n^{-1/2}$) is then similar to Example 11 above. From

$$G^*(s) = \frac{1}{2} \Gamma\left(\frac{s}{2}\right) \zeta^2(s),$$

one gets

$$(42) \quad G(x) \sim \frac{1}{4} \sqrt{\frac{\pi}{x}} (-2 \log(2x) + 3\gamma) + \frac{1}{4} + \dots$$

The other sums $S_n^{(\pm 1)}$ are treated similarly. From (40), (41), (42) (and their analogues), it is found that *the expected height of a random plane rooted tree of n nodes is*

$$\sqrt{\pi n} - \frac{1}{2} + o(1).$$

Full asymptotic expansions could also be determined by this technique.

The basic method here consists in approximating Catalan sums (36) by Gaussian sums of the form

$$G(x) = \sum_{k=1}^{\infty} \lambda_k e^{-k^2 x^2},$$

and treating the latter by Mellin transforms. Related Catalan sums surface in the analysis of Batchier's odd-even merge sorting network [75] and in register allocation [30, 50], where the arithmetic function λ_k involved in (36) is either a function of the Gray code representation of k or the function $v_2(k)$ representing the exponent of 2 in the prime number decomposition of k .

6. THE BERNOULLI SPLITTING PROCESS AND DYADIC SUMS

The Bernoulli splitting process is a general model of the random allocation of resource either in the time domain (like stations sharing a common communication channel) or in the space domain (like keys sharing some primary or secondary storage) whose analysis usually leads to a variety of dyadic sums. The process takes a set G of individuals and splits them recursively as follows:

- If $\text{card}(G) \leq 1$ then the process stops and no splitting occurs.
- Otherwise $\text{card}(G) \geq 2$, and each $g \in G$ flips a coin. Let G_0 and G_1 be the two subsets of G corresponding to the the groups of individuals having flipped heads (0) and tails (1). Then the process is recursively applied to the two subsets G_0 and G_1 .

A realisation of the process may be described by a tree $\tau(G)$ whose internal binary nodes correspond to splittings of more than 1 element; the external nodes either contain single individual or the empty set.

If one views the elements of G as having predetermined an infinite sequence of bits, the tree $\tau(G)$ is nothing but the *digital trie* associated to G viewed as a set of "keys", see [37, 56, 62, 76]. Retrieval of an element g in $\tau(G)$ is achieved by following an access path dictated by g . A sequential execution of the splitting process also constitutes a way to regulate access to a common shared channel (groups consisting of single individuals may deliver their message without interference): this is the *tree communication protocol* of Capetanakis-Tsybakov, see [60, 63].

Given that the cardinality of the original group G is n , there are two basic random variables: the number I_n of nontrivial separation stages corresponding to the number of internal nodes in $\tau(G)$; the total number L_n of coin flippings corresponding to an internal path length in $\tau(G)$. The expectations $i_n = E\{I_n\}$ and $\ell_n = E\{L_n\}$ satisfy recurrences that reflect the nature of the splitting process; for $n \geq 2$, one has

$$(43) \quad i_n = 1 + \sum_{k=0}^n \pi_{n,k}(i_k + i_{n-k}), \quad \ell_n = n + \sum_{k=0}^n \pi_{n,k}(\ell_k + \ell_{n-k}), \quad \pi_{n,k} = \frac{1}{2^n} \binom{n}{k},$$

with initial conditions $i_0 = i_1 = \ell_0 = \ell_1$. The splitting probabilities $\pi_{n,k}$ are specific of the Bernoulli splitting process and they represent the probability of turning k heads out of n coin flips.

The basic technique to solve (43) is to introduce the exponential generating functions

$$(44) \quad I(z) = \sum_{n=0}^{\infty} i_n \frac{z^n}{n!}, \quad L(z) = \sum_{n=0}^{\infty} \ell_n \frac{z^n}{n!},$$

with which (43) transforms into

$$(45) \quad I(z) = 2e^{z/2}I\left(\frac{z}{2}\right) + (e^z - 1 - z), \quad L(z) = 2e^{z/2}L\left(\frac{z}{2}\right) + z(e^z - 1).$$

A functional equation of the form

$$(46) \quad \phi(z) = 2e^{z/2}\phi\left(\frac{z}{2}\right) + a(z),$$

with $a(z)$ a known function and $\phi(z)$ the unknown, is solved by iteration:

$$(47) \quad \begin{aligned} \phi(z) &= a(z) + 2e^{z/2}\phi\left(\frac{z}{2}\right) \\ &= a(z) + 2e^{z/2}a\left(\frac{z}{2}\right) + 4e^{3z/4}a\left(\frac{z}{4}\right) \\ &= \dots \\ &= \sum_{k=0}^{\infty} 2^k e^{z(1-2^{-k})} a\left(\frac{z}{2^k}\right). \end{aligned}$$

This principle applies to $I(z)$ and $L(z)$ with $a(z) = e^z - 1 - z$ and $a(z) = z(e^z - 1)$, respectively. Upon expanding the exponentials, one finds the explicit forms

$$(48) \quad i_n = \sum_{k=0}^{\infty} 2^k \left[1 - \left(1 - \frac{1}{2^k}\right)^n - \frac{n}{2^k} \left(1 - \frac{1}{2^k}\right)^{n-1} \right], \quad \ell_n = n \sum_{k=0}^{\infty} \left[1 - \left(1 - \frac{1}{2^k}\right)^{n-1} \right].$$

From there, the most direct route is the exponential approximation

$$(1 - a)^n = e^{-n \log(1-a)} = e^{-na + \mathcal{O}(na^2)} \approx e^{-na}.$$

It is legitimate to use it in (48) (see [56, p. 131] for a justification based on splitting the sum). With

$$F(x) = \sum_{k=0}^{\infty} 2^k \left[1 - \left(1 + \frac{x}{2^k}\right) e^{-x/2^k} \right] \quad G(x) = x \sum_{k=0}^{\infty} \left[1 - e^{-x/2^k} \right],$$

one finds elementarily $i_n = F(n) + \mathcal{O}(\sqrt{n})$ and $\ell_n = G(n) + \mathcal{O}(\sqrt{n})$. The functions $F(x)$ and $G(x)$ are dyadic sums of a type already considered. Hence

$$\begin{aligned} F(x) &= \frac{x}{\log 2} + xP(\log_2 x) + \mathcal{O}(\sqrt{x}) \\ G(x) &= x \log_2 x + \left(\frac{\gamma}{\log 2} + \frac{1}{2} \right) x + xQ(\log_2 x) + \mathcal{O}(\sqrt{x}), \end{aligned}$$

where $P(u)$ and $Q(u)$ are absolutely convergent Fourier series.

Returning to tries and neglecting the periodic fluctuations that are of amplitude less than 10^{-5} , we find that the number of binary nodes is on average about $1.44n$, a 44% waste in storage, while the average depth ℓ_n/n of a random external node is about $\log_2 n$ which, in the information-theoretic sense, corresponds to an asymptotically optimal search cost.

The variances of these parameters have been extensively studied by Kirschenhofer, Prodinger, and Szpankowski [53, 54]. These authors note that it is not a trivial problem to obtain the exact order of the dominant terms as there are non-trivial cancellations equivalent to modular type identities discovered by Ramanujan.

Jacquet and Régnier [45] have established that path length and node size are both asymptotically normally distributed. The analysis involves the study of nonlinear bivariate difference equations of which the following is typical

$$T(u, z) = uT^2(u, \frac{z}{2}) + (1 - u)(1 + z)e^{-z}.$$

Information for large complex z with u taken as a parameter can be gathered by a quasi-linearization process based on considering $\log T(u, z)$ which is amenable to Mellin transform techniques. Once this is done, the coefficient $[z^n]T(u, z)$ can be recovered asymptotically, and since it is directly related to a characteristic function of the number of nodes, the Gaussian result is then established. Mahmoud's book [62] contains a detailed description of this interesting application of Mellin transforms to nonlinear bivariate problems. A recent extension has been made to suffix trees and the Lempel-Ziv data compression scheme [47].

7. OTHER COMBINATORIAL EXAMPLES

Combinatorial sums may necessitate a certain amount of "preprocessing" before being reducible to harmonic sums, as already exemplified by Catalan sums or sums related to tries. This section describes indirect applications of the asymptotic analysis of harmonic sums to two new types of combinatorial problems.

Reduction to standard harmonic sums. Longest runs in random binary strings are treated by Knuth [57] in a paper that deals with the equivalent problem of carry propagation in parallel binary adders. There the problem requires an analysis of dominant poles of a family of rational functions eventually leading to dyadic sums.

EXAMPLE 16. Longest runs in strings. Consider strings over a binary alphabet $\mathcal{A} = \{0, 1\}$. The problem is to estimate the expected length \bar{L}_n of the longest run of 1's in a random string of length n , where all the 2^n possible strings are taken equally likely. The distribution was studied by Feller [21] and Knuth [57].

The probability that a random string of length n has no run of k consecutive 1's is

$$(49) \quad q_{n,k} = \frac{1}{2^n} [z^n] \frac{1 - z^k}{1 - 2z + z^{k+1}}.$$

The set of such strings is described by the regular expression $1^{<k} \cdot (01^{<k})^*$, where $1^{<k}$ denotes a sequence of less than k 1's and $()^*$ denotes arbitrary repetition of a pattern; the general principles of combinatorial analysis permit to write the ordinary generating function of the set of strings under consideration as

$$\frac{1 - z^k}{1 - z} \cdot \frac{1}{1 - z \frac{1 - z^k}{1 - z}},$$

which justifies (49).

Let ρ_k be the smallest positive root of the denominator of (49) that lies between $\frac{1}{2}$ and 1. An application of the principle of the argument shows such a root to exist with all other

roots that are of a larger modulus. By dominant pole analysis, the $q_{n,k}$ satisfy

$$(50) \quad q_{n,k} \sim c_k(2\rho_k)^{-n} \quad \text{with} \quad c_k = \frac{1 - \rho_k^k}{\rho_k(2 - (k+1)\rho_k^k)},$$

for large n but fixed k .

The denominator of the fraction in (49) behaves near $z = 1/2$ like a “perturbation” of $1 - 2z$ so that one expects ρ_k to be approximated by $\frac{1}{2}$ as $k \rightarrow \infty$. An elementary argument detailed in [57] shows that in fact

$$(51) \quad \rho_k = \frac{1}{2} (1 + 2^{-k-1} + \mathcal{O}(k2^{-2k})).$$

Accordingly $c_k = 1 + \mathcal{O}(k2^{-k})$.

By means of contour integration, Knuth justifies the use of (51) inside (50) for a wide range of values of k and n , which results in the approximate formula:

$$q_{n,k} \approx (1 - 2^{-k-1})^n \approx e^{-n2^{-k-1}}.$$

Let $\hat{q}_{n,k}$ denote the approximation $e^{-n2^{-k-1}}$ to $q_{n,k}$. Following [57], one finds

$$\bar{L}_n \equiv \sum_{k=0}^{\infty} [1 - q_{n,k}] = \sum_{k=0}^{\infty} [1 - \hat{q}_{n,k}] + \mathcal{O}\left(\frac{1}{\sqrt{n}}\right) = \sum_{k=0}^{\infty} [1 - e^{-n2^{-k-1}}] + \mathcal{O}\left(\frac{1}{\sqrt{n}}\right).$$

This is a typical case of a dyadic sum studied repeatedly in previous sections and

$$\bar{L}_n = \log_2 n + \frac{\gamma}{\log 2} - \frac{1}{2} + \frac{1}{\log 2} \sum_{k \in \mathbb{Z} \setminus \{0\}} \Gamma\left(\frac{2ik\pi}{\log 2}\right) e^{-2ik\pi \log_2 n} + \mathcal{O}\left(\frac{1}{\sqrt{n}}\right).$$

Thus, the expected length of the longest run \bar{L}_n fluctuates around $\log_2 n + 0.33274$ with a minute amplitude. \square

An entirely similar analysis provides the expected size of the largest summand in a random composition of an integer n .

Nonstandard Dirichlet series. The algorithm of *Probabilistic Counting* introduced in [27] permits to estimate within a few percents the number of distinct elements of a large file using only a very small amount of auxiliary memory and is of interest in the context of query optimization in data bases. The analysis appeals to the nonstandard Dirichlet series

$$N(s) = \sum_{j=1}^{\infty} \frac{(-1)^{\nu(j)}}{j^s},$$

where $\nu(j)$ is the sum of the digits in the binary representation of j .

EXAMPLE 17. Probabilistic counting. The very design of the algorithm of [27] necessitates the analysis of a parameter of \mathcal{B}^n where $\mathcal{B} = \{0, 1\}^\infty$ is the set of all infinitely long binary strings. Let $u = (u_1, \dots, u_n)$ be an element of \mathcal{B}^n ; the parameter $R = R_n$ is defined to take the value k if the sequences $1, 01, 0^21, \dots, 0^{k-1}1$ all occur as initial segments of some of the u_j 's but no u_j starts with the pattern 0^k1 . A plausible reasoning suggests that the expectation \bar{R}_n of R over \mathcal{B}^n taken with uniform measure should be close to $\log_2 n$, but higher order asymptotic information is required here.

First, an inclusion-exclusion argument of [27] shows that

$$(52) \quad q_{n,k} \equiv \Pr\{R_n \geq k\} = \sum_{j=0}^{2^k} (-1)^{\nu(j)} \left(1 - \frac{j}{2^k}\right)^n.$$

It is proved in [27] that, in a central region near $\log_2 n$ one can use the exponential approximation $(1-a)^n \approx e^{-na}$ while simultaneously extending the range of values of k to infinity. This justifies the approximation

$$(53) \quad q_{n,k} \approx \hat{q}_{n,k} \quad \text{where} \quad \hat{q}_{n,k} = \sum_{j=0}^{\infty} (-1)^{\nu(j)} e^{-jn/2^k} = \prod_{j=0}^{\infty} (1 - e^{-2^j(n/2^k)}).$$

Define the function

$$(54) \quad \theta(x) = \sum_{j \geq 0} (-1)^{\nu(j)} e^{-jx} = \prod_{j \geq 0} (1 - e^{-x2^j}).$$

Eq. (53) means that the cumulative distribution $q_{n,k}$ is well approximated by $\theta(n/2^k)$. In fact, an elementary argument of [27] establishes that

$$(55) \quad \bar{R}_n \equiv \sum_{k=1}^{\infty} q_{n,k} = \Theta(n) + o(1) \quad \text{where} \quad \Theta(x) = \sum_{k=1}^{\infty} \theta\left(\frac{x}{2^k}\right),$$

and the asymptotic form of $\Theta(x)$ for large x is required.

The functions $\theta(x)$ and $\Theta(x)$ are both harmonic sums. From (54) and (55), one finds formally

$$(56) \quad \theta^*(s) = N(s) \cdot \Gamma(s), \quad \Theta^*(s) = \frac{2^s \theta(s)}{1 - 2^s} = \frac{2^s}{1 - 2^s} N(s) \Gamma(s).$$

In particular, a more detailed investigation of $N(s)$ is needed.

Grouping terms by 2^m in the definition of $N(s)$ and using the binomial theorem (see also Prop. 6 below) shows that $N(s)$ is an entire function and is of moderate growth in any right half-plane, owing to cancellations afforded by the sign alternation properties of the sequence $(-1)^{\nu(j)}$. This permits to justify (55) and hence by a residue computation it gives the asymptotic form of $\Theta(x)$:

$$(57) \quad \Theta(x) \underset{x \rightarrow +\infty}{=} \log_2 x + \frac{\gamma}{\log 2} + \frac{N'(0)}{\log 2} - \frac{1}{2} + P(\log_2 x) + o(1),$$

for some oscillating function $P(u)$ once more found to be of small amplitude.

The grouping technique yields $N(0) = -1$ (used in (57)) and it permits to express $N'(0)$ as an alternating sum of logarithms. Thus, from (55), (57), one gets

$$R_n = \log_2(\varphi n) + P(\log_2 n) + o(1),$$

where

$$\varphi = 2^{-1/2} e^{\gamma} \prod_{m=1}^{\infty} \left(\frac{2m+1}{2m} \right)^{(-1)^{\nu(m)}} \doteq 0.77351.$$

There results an algorithm based on the idea that $\frac{1}{\varphi} 2^{R_n}$ can be turned into an unbiased statistical estimator of the *a priori* unknown number n of distinct binary strings considered.

□

The sequence $\varepsilon(j) = (-1)^{\nu(j)}$ is the classical Thue-Morse sequence. Dirichlet series related to $N(s)$ have been considered in [2]. Similar techniques are used in other probabilistic estimation algorithms like *Approximate Counting* [22] and the collision resolution methods of [39].

III. General Mellin Asymptotics

The last sections have shown that Mellin asymptotics is applicable to harmonic sums provided the Dirichlet series is of moderate growth while the transform of the base function is of fast decrease. The applications encountered so far have made an essential use of properties specific to the Riemann zeta function, to the Gamma function, or to the sine function.

This part shows that, thanks to general theorems presented in Section 8, Mellin asymptotics may also be applied to “implicit” harmonic sums, where it is no longer required to have closed forms for either the Dirichlet series or the transform of the base function.

8. GENERAL CONDITIONS FOR MELLIN ASYMPTOTICS

In this section, we provide general conditions under which Mellin transforms are small towards $\pm i\infty$ and describe a general class of Dirichlet series that are of moderate growth. These results extend the range of applicability of Mellin asymptotics to a much larger class of harmonic sums.

Smallness of Mellin transforms. The smallness of a Mellin transform is directly related to the degree of “smoothness” (differentiability, analyticity) of the original function.

Proposition 3 (Smallness in the fundamental strip). (i) *Let $f(x)$ be locally integrable with fundamental strip (α, β) . Then, uniformly with respect to σ in any closed subinterval of (α, β) , one has*

$$f^*(\sigma + it) = o(1) \quad \text{as } t \rightarrow \pm\infty.$$

(ii) *If in addition $f(x)$ is of class C^r and the fundamental strip of $\Theta^r f$ contains (α, β) , then*

$$f^*(\sigma + it) = o(|t|^{-r}) \quad \text{as } t \rightarrow \pm\infty.$$

Proof. (Sketch) (i) From the form

$$f^*(\sigma + it) = \int_0^\infty f(x) e^{-\sigma x} e^{it \log x} dx,$$

the function $f^*(s)$ is an integrable function hashed by a complex exponential. By the Riemann-Lebesgue lemma [43, 81], $f^*(s)$ tends to 0 as $t \rightarrow \pm\infty$.

(ii) The second form results from the formula $\mathcal{M}[\Theta^r f(x); s] = (-1)^r s^r f^*(s)$. \square

The next proposition shows that smallness extends beyond the fundamental strip for smooth functions with smooth derivatives.

Proposition 4 (Smallness beyond the fundamental strip). *Let $f(x)$ be of class C^r with fundamental strip $\langle \alpha, \beta \rangle$. Assume that $f(x)$ admits an asymptotic expansion as $x \rightarrow 0^+$ (resp. $x \rightarrow +\infty$) of the form*

$$(58) \quad f(x) = \sum_{(\xi, k) \in A} c_{\xi, k} x^{\xi} (\log x)^k + \mathcal{O}(x^{\gamma})$$

where the ξ satisfy $-\alpha \leq \xi < \gamma$ (resp. $\gamma < \xi \leq -\beta$). Assume also that each derivative $\frac{d^j}{dx^j} f(x)$ for $j = 1, \dots, r$ satisfies an asymptotic expansion obtained by termwise differentiation of (58). Then the continuation of $f^*(s)$ satisfies

$$(59) \quad f^*(\sigma + it) = o(|t|^{-r}) \quad \text{as } |t| \rightarrow \infty$$

uniformly for σ in any closed subinterval of $(-\gamma, \beta)$ (resp. of $(\alpha, -\gamma)$).

Proof. It suffices to consider extension to the left of $f^*(s)$. Choose some positive number $p > \gamma$ and define

$$a(x) = \left(\sum_{(\xi, k) \in A} c_{\xi, k} x^{\xi} (\log x)^k \right) \exp(-x^p).$$

The function $g(x) = f(x) - a(x)$ satisfies the assumptions of Proposition 3 and its transform $g^*(s)$ is thus $o(|s|^{-r})$ in its fundamental strip $\langle -\gamma, \beta \rangle$. The transform $a^*(s)$ is itself exponentially small given growth properties of the Gamma function and its derivatives. Thus $f^*(s) = a^*(s) + g^*(s)$ satisfies the stated bounds. \square

Analyticity is the strongest possible form of smoothness for a function $f(x)$; in that case the transform $f^*(s)$ decays exponentially in a quantifiable way.

Proposition 5 (Exponential smallness in the analytic case). *Let $f(x)$ be analytic in S_{θ} where S_{θ} is the sector*

$$S_{\theta} = \{z \in \mathbb{C} \mid 0 < |t| < +\infty \text{ and } |\arg(z)| \leq \theta\} \quad \text{with } 0 < \theta < \pi.$$

Assume that $f(x) = \mathcal{O}(x^{-\alpha})$ as $x \rightarrow 0$ in S_{θ} , and $f(x) = \mathcal{O}(x^{-\beta})$ as $x \rightarrow \infty$ in S_{θ} . Then,

$$f^*(\sigma + it) = \mathcal{O}(e^{-\theta|t|})$$

uniformly for σ in every closed subinterval of (α, β) .

Proof. (Sketch) The integral defining Mellin transforms is in this case applied to an analytic function. By Cauchy's theorem, the integration contour may be taken as the half-line of slope θ :

$$f^*(s) = \int_0^{e^{i\theta}\infty} f(t) t^{s-1} dt.$$

The change of variable $t = \rho e^{i\theta}$ gives

$$f^*(s) = e^{i\theta s} \int_0^{\infty} f(\rho e^{i\theta}) \rho^{s-1} d\rho.$$

The result follows as the integral converges. \square

Smallness extends outside of the fundamental strip by an argument similar to that of Prop. 4 and based on subtracting combinations of exponentials.

EXAMPLE 18. *A sum with an implicit transform.* The harmonic sum

$$G(x) = \sum_{n=1}^{\infty} \frac{d(n)}{\sqrt{\cosh nx}},$$

with $d(n)$ the divisor function, admits the transform $\zeta^2(s) \cdot g^*(s)$ where $g^*(s)$ is the transform of $g(x) = (\cosh x)^{-1/2}$. The function $g^*(s)$ is not simply expressible in terms of standard functions (it is “implicitly” defined as the transform of $g(x)$). However, the fundamental correspondence provides its complete singular expansion (see Example 4) while Propositions 4 and 5 show that $g^*(s)$ is of fast decrease in any right half-plane. Thus, Mellin analysis applies and the asymptotics of $G(x)$ as $x \rightarrow 0$ obtains from the singular expansions of $g^*(s)$ and $\zeta^2(s)$. One gets:

$$G(x) \underset{x \rightarrow 0}{\sim} -g^*(1) \frac{\log x}{x} + \frac{(g^*)'(1) + 2\gamma g^*(1)}{x} + \frac{1}{4} + \mathcal{O}(x^M),$$

for any $M > 0$. There $g^*(1) = \int_0^\infty g(x) dx$ and $(g^*)'(1)$ is an “Euler constant” of $g(x)$ which is expressible as an integral as discussed in Example 15. \square

EXAMPLE 19. *A lattice sum.* The goal here is to discuss certain (2-dimensional) “lattice sums” of the form

$$(60) \quad F(x) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} f((m^2 + n^2)x^2).$$

(See also [10] for related developments.) Such a sum may be rewritten as $\sum_{k \geq 1} r(k) f(kx^2)$, where $r(k)$ is the number of representations of k as a sum of two squares. The corresponding Dirichlet series³ is then

$$\rho(s) = \sum_{k=1}^{\infty} \frac{r(k)}{k^s} = \sum_{m,n \geq 1} \frac{1}{(m^2 + n^2)^s},$$

and everything rests on its behaviour at $\pm i\infty$ and its special values.

Let $\Theta(x) = \sum_{m=1}^{\infty} e^{-m^2 x^2}$. The Mellin transform of $\Theta^2(x)$ is

$$\mathcal{M}(\Theta^2(x), s) = \rho(s) \cdot \frac{1}{2} \Gamma\left(\frac{s}{2}\right).$$

An already discussed Mellin analysis of $\Theta(x)$ yields its asymptotic expansion near $x = 0$:

$$(61) \quad \Theta(x) = \sqrt{\frac{\pi}{2x}} - \frac{1}{2} + R(x)$$

the remainder term $R(x)$ being exponentially small (this alternatively results from Poisson’s formula). Thus, $\Theta^2(x)$ admits an explicit three-term asymptotic expansion,

$$\Theta^2(x) = \frac{\pi}{2x} - \sqrt{\frac{\pi}{2x}} + \frac{1}{4} + R_2(x),$$

³In order to allow for extensions to higher dimensions or higher powers, we do not make use here of special properties of $\rho(s)$. See [40, Ch. IX] for a vivid account of these aspects.

with again $R_2(x)$ exponentially small.

Consequently, $\rho(s)$ is meromorphic in the whole of \mathbb{C} with poles at $s = 2, 1$ only and singular expansion

$$\rho(s) \asymp \frac{\pi}{2(s-2)} - \frac{1}{s-1} + \left[\frac{1}{4s} \right]_{s=0} + [0]_{s=-2} + [0]_{s=-4} + \dots$$

In addition, Proposition 5 quantifies the growth of the transform of $\Theta^2(x)$ in the form $e^{-(\pi/4-\epsilon)|t|}$. Thus, $\rho(s)$ is itself of growth at most $e^{\epsilon|s|}$ for any $\epsilon > 0$ in any finite strip of the complex plane

This knowledge permits to analyse lattice sums of the form (60). For instance,

$$L(p) = \sum_{m,n \geq 1} \frac{1}{(m^2 + n^2 + p^2)^3}$$

satisfies $L(p) = p^{-6}F(1/p)$ provided one takes $f(x) = (1+x)^{-3}$. One has in $(0, 3)$

$$f^*(s) = \frac{\pi}{\sin \pi s} \frac{(s-1)(s-2)}{2}.$$

There results an asymptotic form of $F(x)$ that induces a corresponding estimate of $L(p)$:

$$L(p) \sim \frac{\pi}{8p^4} - \frac{3\pi}{16p^5} + \frac{5-\pi}{4p^6},$$

the error term being exponentially small since $\rho(s)$ vanishes at the even negative integers. The method clearly extends to higher dimensional sums. \square

Growth of Dirichlet series. Dirichlet series whose components λ_k, μ_k admit descending asymptotic expansions in powers of k have the property of being meromorphically continuable with well individuated poles.

Proposition 6 (Growth of special Dirichlet series). *Let λ_k and μ_k admit asymptotic expansions in descending powers of k as $k \rightarrow \infty$:*

$$\lambda_k \sim \sum_{r=0}^{\infty} \frac{a_r}{k^{\alpha_r}}, \quad \mu_k \sim k^w \left(1 + \sum_{r=1}^{\infty} \frac{b_r}{k^{\beta_r}} \right).$$

Then the Dirichlet series $\sum_k \lambda_k \mu_k^{-s}$ can be continued to a meromorphic function $\Lambda(s)$ in the whole of the complex plane. The function $\Lambda(s)$ is of moderate growth in any right half plane of the complex plane.

Proof. (Sketch) The particular case $\alpha_r = \beta_r = r$ gives the essential idea. The binomial theorem yields

(62)

$$\begin{aligned} \frac{\lambda_n}{\mu_n^s} &= \frac{1}{n^{ws}} \left[a_0 + \frac{a_1}{n} + \dots \right] \left[1 - \frac{s}{1!} \left(\frac{b_1}{n} + \dots \right) + \frac{s(s+1)}{2!} \left(\frac{b_1}{n} + \dots \right)^2 + \dots \right] \\ &= \frac{a_0}{n^{ws}} + \sum_{k \geq 1} \frac{P_k(s)}{n^{ws+k}}, \end{aligned}$$

for a family of polynomials P_k . By summation, one finds that $\Lambda(s)$ has, in any finite strip, the same singularities as a partial sum of

$$\zeta(ws) + \sum_{k \geq 1} P_k(s) \zeta(ws + k).$$

(This is easily justified rigorously by terminating forms of the binomial theorem.) Also, $\Lambda(s)$ is of moderate growth since each component zeta function shares this characteristic.

In the general case, one finds similarly (with a slight abuse of notations)

$$(63) \quad \Lambda(s) \asymp \zeta(ws + \alpha_0) + \sum_{\xi \in \Xi} P_\xi(s) \zeta(ws + \xi),$$

where ξ ranges over the set Ξ of all linear combinations of the form $\alpha_{r_0} + \beta_{r_1} + \cdots + \beta_{r_j}$. \square

For instance, from the expansion

$$\frac{1}{4^n} \binom{2n}{n} = \frac{1}{\sqrt{\pi n}} \left[1 - \frac{1}{8n} + \frac{1}{128n^2} + \mathcal{O}\left(\frac{1}{n^3}\right) \right],$$

one finds by summation that

$$(64) \quad \Lambda(s) \equiv \sum_{n=1}^{\infty} \frac{1}{4^n} \binom{2n}{n} \frac{1}{n^s} = \frac{1}{\sqrt{\pi}} \left[\zeta\left(s + \frac{1}{2}\right) - \frac{1}{8} \zeta\left(s + \frac{3}{2}\right) + \frac{1}{128} \zeta\left(s + \frac{5}{2}\right) + R(s) \right],$$

where $R(s)$ converges in the same half plane as $\zeta\left(s + \frac{7}{2}\right)$, namely in $\langle -\frac{5}{2}, +\infty \rangle$.

In a similar vein, if $Q(u)$ is a polynomial that does not vanish at the positive integers, and $P(u)$ is an arbitrary polynomial, then

$$\Lambda(s) = \sum_{n=1}^{\infty} \frac{P(n)}{(Q(n))^s}$$

has poles only at a discrete set of rational numbers that is bounded from the right.

Amplitudes involving the harmonic numbers $H_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n}$ can be treated in this way by considering $\hat{H}_n = H_n - \log n$. From

$$H_n = \log n + \gamma + \frac{1}{2n} - \frac{1}{12n^2} + \frac{1}{120n^4} + \mathcal{O}(n^{-6}),$$

one deduces

$$\Lambda(s) \equiv \sum_{n=1}^{\infty} \frac{H_n}{n^s} \asymp \zeta'(s) + \gamma \zeta(s) + \frac{1}{2} \zeta(s+1) - \frac{1}{12} \zeta(s+2) + \frac{1}{120} \zeta(s+4) \quad (s \in \langle -5, \infty \rangle).$$

For instance, $\Lambda(s)$ has a simple pole with residue $1/120$ at $s = -3$. More information on this function may be found in [5].

EXAMPLE 20. *A sum with binomial amplitudes.* The function

$$G(x) = \left(\frac{1}{2}\right) \frac{1}{1+x} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right) \frac{1}{1+2x} + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right) \frac{1}{1+3x} + \dots$$

has the transform $\Lambda(s)\pi/\sin \pi s$ where $\Lambda(s) = \sum_{n=1}^{\infty} 4^{-n} \binom{2n}{n} n^{-s}$. The singularities of $\Lambda(s)$ are given by Eq. (64), hence

$$G(x) \underset{x \rightarrow 0}{=} \sqrt{\frac{\pi}{x}} + \Lambda(0) + \frac{1}{8} \sqrt{\pi x} + \mathcal{O}(x),$$

where $\Lambda(0)$ is obtained by the analytic continuation technique of Proposition 6:

$$\Lambda(0) = \sum_{n=1}^{\infty} \left[\frac{1}{4^n} \binom{2n}{n} - \frac{1}{\sqrt{\pi n}} \right] + \frac{1}{\sqrt{\pi}} \zeta\left(\frac{1}{2}\right).$$

A full asymptotic expansion in powers of $x^{1/2}$ can be determined in this way. \square

EXAMPLE 21. *Stirling's formula for modified Gamma functions.* Consider the problem of evaluating asymptotically as $x \rightarrow +\infty$

$$H(x) = \prod_{n=1}^{\infty} \left(1 + \frac{x}{n(n+1)}\right).$$

One estimates instead the harmonic sum $G(x) = \log H(x)$ that is a harmonic sum with transform valid in $\langle -1, -\frac{1}{2} \rangle$:

$$G^*(s) = \Lambda(-s) \cdot \left(\frac{\pi}{s \sin \pi s}\right) \quad \text{where} \quad \Lambda(s) = \sum_{n=1}^{\infty} \frac{1}{(n^2 + n)^s}.$$

From Proposition 6, $\Lambda(s)$ has a singular expansion induced by the expansion of $(1+n^{-1})^{-s}$. One has

$$\Lambda(s) = \zeta(2s) - \frac{s}{1} \zeta(2s+1) + R(s)$$

where $R(s)$ converges like $\zeta(2s+2)$ in $\Re(s) > -\frac{1}{2}$.

The Mellin transform $G^*(s)$ thus has a simple pole at $s = -\frac{1}{2}$ and a triple pole at 0. This provides "Stirling's formula" for $H(x)$:

$$H(x) \underset{x \rightarrow \infty}{\sim} e^{\pi\sqrt{x}} \frac{C}{x}$$

where "Stirling's constant" C is here $C = 1/\pi$.

In this particular case, $H(x)$ is explicitly expressible (by the infinite product formula for $\cos(x)$); the example is only meant to demonstrate the general approach to the analysis of such modified Gamma functions. \square

A similar technique works for Dirichlet series $\Lambda(s)$ whose elements λ_k, μ_k admit expansions in powers of 2^k . Assume that

$$\lambda_k \sim 1 + \sum_{j=1}^{\infty} a_j 2^{-kj}, \quad \mu_k \sim 2^k \left(1 + \sum_{j=1}^{\infty} b_j 2^{-kj}\right).$$

A technique entirely similar to that of Proposition 6 applies. From

$$\lambda_k(\mu_k)^{-s} \sim 2^{-ks} \cdot \left(1 + (a_1 - b_1 s)2^{-k} + (a_2 + \frac{1}{2}(b_1^2 - 2a_1 b_1 - 2b_2)s + \frac{1}{2}b_1^2 s^2)2^{-2k} + \dots \right),$$

one gets by summation

$$\Lambda(s) = \frac{2^{-s}}{1 - 2^{-s}} + (a_1 - b_1 s) \frac{2^{-s-1}}{1 - 2^{-s-1}} + \dots.$$

Thus, $\Lambda(s)$ admits a pole at all points of a half-lattice

$$\chi_k^{(m)} = -m + \frac{2ik\pi}{\log 2}, \quad \text{for } m \geq 0, k \in \mathbb{Z},$$

and is in addition of controlled growth away from poles in any finite strip of \mathbb{C} .

An immediate case of application is to modified dyadic sums involving frequencies of the form $2^k \pm 1$. This technique also applies to sums related to the Bernoulli splitting model of Section 6; for instance, the Mellin transform of

$$F(x) = \sum_{k=1}^{\infty} \left[1 - \left(1 - \frac{1}{2^k} \right)^x \right]$$

is

$$F^*(s) = \Lambda(s) \cdot \Gamma(s) \quad \text{where} \quad \Lambda(s) = \sum_{k=1}^{\infty} (\log(1 - 2^{-k})^{-1})^{-s}.$$

By the technique above one finds

$$\Lambda(s) \asymp \frac{2^s}{1 - 2^s} - \frac{s}{2} \frac{2^{s-1}}{1 - 2^{s-1}} \quad (s \in \langle -\infty, 2 \rangle).$$

This permits to compute complete asymptotic expansions without a recourse to the exponential approximation. The half-lattice of poles implies the presence of fluctuating functions at each level of the asymptotic expansion of $F(x)$.

The transfer technique. Our final example illustrates the general approach seen so far in conjunction with an important transfer technique that consists in going back and forth between various harmonic sums involving related amplitude and frequencies but different base functions. This “zigzag” method appeals to the common occurrence of a Dirichlet series in two harmonic sums

$$\sum a_k f(kx) \quad \text{and} \quad \sum a_k g(kx),$$

with $g(x) = e^{-x}$ playing a special rôle because of the relation it establishes with ordinary generating functions.

The transfer technique has already been used implicitly: for instance the study of the harmonic sum $e^{-x}(1 - e^{-x})^{-1}$ yields properties of $\zeta(s)$ (Example 2) that can in turn be used to analyze the theta function $\sum_k e^{-k^2 x^2}$ (Example 10) and eventually lattice sums (Example 19).

The principle is as follows. Assume that the behaviour of

$$F(x) = \sum_{k=1}^{\infty} a_k f(kx)$$

is sought. This requires knowledge of the Dirichlet series

$$\alpha(s) = \sum_{k=1}^{\infty} \frac{a_k}{k^s}.$$

If the ordinary generating function

$$A(x) = \sum_{k=1}^{\infty} a_k x^k$$

is known and in addition $A(x)$ is analytic at $x = 1$ (this often happens for alternating series with elementary coefficients), then

$$A(e^{-x}) = \sum_{m=0}^{\infty} b_m x^m,$$

which corresponds to a singular expansion

$$\mathcal{M}(A(e^{-x}), s) = \alpha(s)\Gamma(s) \asymp \sum_{m=0}^{\infty} \frac{b_m}{m+s} \quad (s \in \mathbb{C}).$$

Comparison with the singular expansion of the product $\alpha(s)\Gamma(s)$, namely

$$\alpha(s) \asymp \sum_{m=0}^{\infty} (-1)^m \frac{\alpha(-m)}{m!} \frac{1}{s+m}$$

shows by identification that

$$(65) \quad \alpha(-m) = (-1)^m m! b_m = (-1)^m m! [x^m] A(e^{-x}).$$

In this way, coefficients in many expansions can be determined explicitly. In general, the formulæ obtained are nontrivial⁴ since the original series defining $\alpha(s)$ stops being convergent at negative integers.

EXAMPLE 22. *An alternating sum with harmonic numbers.* Define

$$G(x) = \sum_{k=1}^{\infty} (-1)^k H_k e^{-k^2 x^2},$$

that arises as an approximation to the combinatorial sum

$$\sum_{k=1}^n (-1)^k H_k \binom{2n}{n-k}.$$

⁴The transfer process often provides a rigorous counterpart of formal computations with purely divergent series. See the remarks relative to Euler-Maclaurin summations in Section 4 and the theory of “regularized” sums and products in [49].

The Mellin transform of $G(x)$ is

$$G^*(s) = h(s) \cdot \frac{1}{2} \Gamma\left(\frac{s}{2}\right) \quad \text{where} \quad h(s) = \sum_{k=1}^{\infty} (-1)^k \frac{H_k}{k^s}.$$

Thus, $G^*(s)$ has poles at $s = 0, -2, -4, \dots$ and determining the residues of $G^*(s)$ requires the values of $h(s)$ at the negative even integers.

The Dirichlet series $h(s)$ is an entire function. This results from a simple extension of Proposition 6 to alternating series, as alternating zeta functions that are entire. The special values of $h(s)$ are then obtained from the expansion

$$\begin{aligned} \sum_{k=1}^{\infty} (-1)^k H_k e^{-kx} &= \frac{1}{1+e^{-x}} \log \frac{1}{1+e^{-x}} \\ &= -\frac{\log 2}{2} + \left(\frac{1}{4} - \frac{\log 2}{4}\right)x + \frac{1}{16}x^2 + \left(-\frac{1}{32} + \frac{\log 2}{48}\right)x^3 - \frac{1}{128}x^4 + \dots, \end{aligned}$$

so that

$$h(0) = -\frac{1}{2} \log 2, \quad h(-2) = \frac{1}{8}, \quad h(-4) = -\frac{3}{16}, \dots$$

Thus finally

$$G(x) \underset{x \rightarrow 0}{\sim} -\frac{1}{2} \log 2 - \frac{1}{8}x^2 - \frac{3}{32}x^4 + \dots$$

□

9. CONCLUSIONS

This paper has demonstrated the basic technology of Mellin asymptotics of harmonic sums. There, a crucial rôle is played by the separation property and the fundamental correspondence. The method is likely to be applicable as soon as the expressions involved are of an analytic character (in the old sense of the term) affording analytic continuation which is the most important requirement of the method. Combinatorial expressions are thus very natural candidates for Mellin asymptotics. Related techniques in connection with finite differences are explored in [34].

As mentioned in Section 4, Perron's formula belongs to the galaxy of Mellin related techniques. We have not developed this classical aspect here as it is extensively discussed elsewhere. In [25] it is shown how divide-and-conquer recurrences lead to fluctuations of a fractal nature that can be quantified via a Mellin analysis; digital sequences are studied along similar lines in [26].

An interesting offspring of Mellin asymptotics of harmonic sums is the class of *two-stage* methods. In many cases, a generating function can be analyzed asymptotically as a harmonic sum in the complex realm, near a singularity or a critical point. It then becomes possible to estimate the coefficients of the generating function by either singularity analysis or the saddle-point method. For a conjunction of Mellin and saddle-point methods, the standard example is De Bruijn's pioneering work on the enumeration of binary partitions [14] which was followed by the general approach of Meinardus in the theory of integer partitions [3]. Singularity analysis used in conjunction with Mellin asymptotics may be used as a basis for the analysis

of Catalan sums [28] as well as many other combinatorial sums. We hope to return to these questions in a future paper.

Acknowledgement. This work was partly supported by the ESPRIT Basic Research Action of the E. C. No. 7141 (ALCOM II).

The authors would like to thank Rainer Kemp (who introduced one of us to the subject in 1978), Robert Sedgewick (for a long lasting interaction), Leo Guibas and Andrew Odlyzko (for numerous discussions), Don Knuth and N. G. de Bruin (for their support). Helmut Prodinger together with his group of collaborators at the Technical University of Vienna, Mireille Régnier, Philippe Jacquet, and Wojtek Szpankowski all contributed valuable feedback at various stages. We are deeply grateful to them for their cooperation.

REFERENCES

1. ABRAMOWITZ, M., AND STEGUN, I. A. *Handbook of Mathematical Functions*. Dover, 1973. A reprint of the tenth National Bureau of Standards edition, 1964.
2. ALLOUCHE, J.-P., AND COHEN, H. Dirichlet series and curious infinite products. *Bulletin of the London Mathematical Society* 17 (1985), 531–538.
3. ANDREWS, G. E. *The Theory of Partitions*, vol. 2 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley, 1976.
4. APOSTOL, T. M. *Introduction to Analytic Number Theory*. Springer-Verlag, 1976.
5. APOSTOL, T. M., AND VU, T. H. Dirichlet series related to the Riemann zeta function. *Journal of Number Theory* 19 (1984), 85–102.
6. BARNES, E. W. The Maclaurin sum formula. *Proceedings of the London Mathematical Society* 2, 3 (1905), 253–272.
7. BELLMAN, R. *Analytic Number Theory, An Introduction*. Benjamin-Cummings, Reading, 1980.
8. BERNDT, B. C. *Ramanujan's Notebooks, Part II*. Springer Verlag, 1989.
9. BERNDT, B. C., AND EVANS, R. J. Extensions of asymptotic expansions from Chapter 15 of Ramanujan's second notebook. *Journal für die reine und angewandte Mathematik* 361 (1985), 118–134.
10. BORWEIN, J. M., BORWEIN, P. B., AND SHAIL, R. Analysis of certain lattice sums. *Journal of Mathematical Analysis and Applications* 143, 1 (1989), 126–137.
11. COMTET, L. *Advanced Combinatorics*. Reidel, Dordrecht, 1974.
12. DAVENPORT, H. *Multiplicative Number Theory*, revised by H. L. Montgomery, second ed. Springer-Verlag, New York, 1980.
13. DAVIES, B. *Integral Transforms and Their Applications*. Springer-Verlag, 1978.
14. DE BRUIJN, N. G. On Mahler's partition problem. *Indagationes Math.* 10 (1948), 210–220. Reprinted from Koninkl. Nederl. Akademie Wetenschappen, Ser. A.
15. DE BRUIJN, N. G. *Asymptotic Methods in Analysis*. Dover, 1981. A reprint of the third North Holland edition, 1970 (first edition, 1958).
16. DE BRUIJN, N. G., KNUTH, D. E., AND RICE, S. O. The average height of planted plane trees. In *Graph Theory and Computing* (1972), R. C. Read, Ed., Academic Press, pp. 15–22.
17. DINGLE, R. B. *Asymptotic Expansions: Their Derivation and Interpretation*. Academic Press, London, 1973.
18. DOETSCH, G. *Handbuch der Laplace Transformation, Vol. 1–3*. Birkhäuser Verlag, Basel, 1955.
19. ERDÉLYI, A., MAGNUS, W., OBERHETTINGER, F., AND TRICOMI, F. G. *Tables of Integral Transforms*. McGraw-Hill, 1954.
20. FAYOLLE, G., FLAJOLET, P., AND HOFRI, M. On a functional equation arising in the analysis of a protocol for a multiaccess broadcast channel. *Advances in Applied Probability* 18 (1986), 441–472.
21. FELLER, W. *An Introduction to Probability Theory and its Applications*, third ed., vol. 1. John Wiley, 1968.
22. FLAJOLET, P. Approximate counting: A detailed analysis. *BIT* 25 (1985), 113–134.
23. FLAJOLET, P. On adaptive sampling. *Computing* 34 (1990), 391–400.
24. FLAJOLET, P., AND GOLIN, M. Exact asymptotics of divide-and-conquer recurrences. In *Automata, Languages, and Programming* (1993), S. C. A. Lingas, R. Karlsson, Ed., no. 700 in Lecture Notes in Computer Science, pp. 137–149. Proceedings of the 20th

ICALP Conference, Lund, July 1993.

25. FLAJOLET, P., AND GOLIN, M. Mellin transforms and asymptotics: The mergesort recurrence. *Acta Informatica* 31 (1994). In press.
26. FLAJOLET, P., GRABNER, P., KIRSCHENHOFER, P., PRODINGER, H., AND TICHY, R. Mellin transforms and asymptotics: Digital sums. *Theoretical Computer Science* 123, 2 (1994), 291-314.
27. FLAJOLET, P., AND MARTIN, G. N. Probabilistic counting algorithms for data base applications. *Journal of Computer and System Sciences* 31, 2 (Oct. 1985), 182-209.
28. FLAJOLET, P., AND PRODINGER, H. Register allocation for unary-binary trees. *SIAM Journal on Computing* 15, 3 (Aug. 1986), 629-640.
29. FLAJOLET, P., AND PUECH, C. Partial match retrieval of multidimensional data. *Journal of the ACM* 33, 2 (1986), 371-407.
30. FLAJOLET, P., RAOULT, J.-C., AND VUILLEMIN, J. The number of registers required to evaluate arithmetic expressions. *Theoretical Computer Science* 9 (1979), 99-125.
31. FLAJOLET, P., RÉGNIER, M., AND SEDGEWICK, R. Some uses of the Mellin integral transform in the analysis of algorithms. In *Combinatorial Algorithms on Words* (1985), A. Apostolico and Z. Galil, Eds., vol. 12 of *NATO Advance Science Institute Series. Series F: Computer and Systems Sciences*, Springer Verlag, pp. 241-254. (Invited Lecture).
32. FLAJOLET, P., AND RICHMOND, B. Generalized digital trees and their difference-differential equations. *Random Structures and Algorithms* 3, 3 (1992), 305-320.
33. FLAJOLET, P., SALVY, B., AND ZIMMERMANN, P. Automatic average-case analysis of algorithms. *Theoretical Computer Science* 79, 1 (Feb. 1991), 37-109.
34. FLAJOLET, P., AND SEDGEWICK, R. Mellin transforms and asymptotics: finite differences and Rice's integrals. Research Report 2031, Institut National de Recherche en Informatique et en Automatique, Mar. 1994. 21 pages. To appear in *Theoretical Computer Science*.
35. FREDMAN, M. L., AND KNUTH, D. E. Recurrence relations based on minimization. *Journal of Mathematical Analysis and Applications* 48 (1974), 534-559.
36. GONNET, G. H. Notes on the derivation of asymptotic expressions from summations. *Information Processing Letters* 7, 4 (1978), 165-169.
37. GONNET, G. H., AND BAEZA-YATES, R. *Handbook of Algorithms and Data Structures: in Pascal and C*, second ed. Addison-Wesley, 1991.
38. GOULDEN, I. P., AND JACKSON, D. M. *Combinatorial Enumeration*. John Wiley, New York, 1983.
39. GREENBERG, A. G., FLAJOLET, P., AND LADNER, R. E. Estimating the multiplicities of conflicts to speed their resolution in multiple access channels. *Journal of the ACM* 34, 2 (Apr. 1987), 289-325.
40. HARDY, G. H. *Ramanujan: Twelve Lectures on Subjects Suggested by his Life and Work*, third ed. Chelsea Publishing Company, New-York, 1978. Reprinted and Corrected from the First Edition, Cambridge, 1940.
41. HARDY, G. H., AND RIESZ, M. *The General Theory of Dirichlet's Series*. No. 18 in *Tracts in Mathematics*. Cambridge University Press, 1915.
42. HECKE, E. *Lectures on Dirichlet Series, Modular Functions and Quadratic Forms*. Vandenhoeck and Rupprecht, Göttingen, 1983. (Edited by Bruno Schoenberg.).
43. HENRICI, P. *Applied and Computational Complex Analysis*. John Wiley, New York, 1977. 3 volumes.
44. HOFRI, M. *Probabilistic Analysis of Algorithms*. Springer-Verlag, 1987.

45. JACQUET, P., AND RÉGNIER, M. Trie partitioning process: Limiting distributions. In *CAAP'86* (1986), P. Franchi-Zanetacchi, Ed., vol. 214 of *Lecture Notes in Computer Science*, pp. 196–210. Proceedings of the 11th Colloquium on Trees in Algebra and Programming, Nice France, March 1986.
46. JACQUET, P., AND SZPANKOWSKI, W. Ultimate characterizations of the brurst response of an interval searching algorithm: A study of a functional equation. *SIAM Journal on Computing* 18, 4 (1989), 777–791.
47. JACQUET, P., AND SZPANKOWSKI, W. Asymptotic behavior of the Lempel-Ziv parsing scheme and digital search trees. preprint, 1991.
48. JACQUET, P., AND SZPANKOWSKI, W. Autocorrelation on words and its applications: Analysis of suffix trees by string-ruler approach. *Journal of Combinatorial Theory, Series A* 66, 2 (1994), 237–269.
49. JORGENSEN, J. A., AND LANG, S. *Basic Analysis of Regularized Series and Products*, vol. 1564 of *Lecture Notes in Mathematics*. Springer Verlag, 1993.
50. KEMP, R. The average number of registers needed to evaluate a binary tree optimally. *Acta Informatica* 11, 4 (1979), 363–372.
51. KEMP, R. *Fundamentals of the Average Case Analysis of Particular Algorithms*. Wiley-Teubner, Stuttgart, 1984.
52. KIRSCHENHOFER, P., MARTINEZ, C., AND PRODINGER, H. Analysis of an optimized search algorithm for skip lists, 1994. 21 pages.
53. KIRSCHENHOFER, P., AND PRODINGER, H. On some applications of formulæ of Ramanujan in the analysis of algorithms. *Mathematika* 38 (1991), 14–33.
54. KIRSCHENHOFER, P., PRODINGER, H., AND SZPANKOWSKI, W. On the variance of the external path length in a symmetric digital trie. *Discrete Applied Mathematics* 25 (1989), 129–143.
55. KLUSCH, D. The sampling theorem, Dirichlet series and Hankel transforms. *Journal of Computational and Applied Mathematics* 44 (1992), 261–273.
56. KNUTH, D. E. *The Art of Computer Programming*, vol. 3: Sorting and Searching. Addison-Wesley, 1973.
57. KNUTH, D. E. The average time for carry propagation. *Indagationes Mathematicae* 40 (1978), 238–242.
58. LEWIN, L., Ed. *Structural Properties of Polylogarithms*. American Mathematical Society, 1991.
59. LINDELÖF, E. Robert Hjalmar Mellin. *Advances in Mathematics* 61 (1933), i–vi. H. Mellin's notice and bibliography.
60. LONGO, G., Ed. *Multi-User Communication Systems*, vol. 265 of *CISM Courses and Lecture Notes*. Springer Verlag, 1981.
61. MACFARLANE, G. G. The application of Mellin transforms to the summation of slowly converging series. *Philosophical Magazine* 40 (1949), 188–197.
62. MAHMOUD, H. *Evolution of Random Search Trees*. John Wiley, New York, 1992.
63. MATHYS, P., AND FLAJOLET, P. Q-ary collision resolution algorithms in random access systems with free or blocked channel access. *IEEE Transactions on Information Theory* IT-31, 2 (Mar. 1985), 217–243.
64. MELLIN, H. Die Dirichlet'schen Reihen, die zahlentheoretischen Funktionen und die endlichen Produkte von endlichem Geschlecht. *Acta Mathematica* 28 (1904), 37–64.
65. MELLIN, H. Abriß einer einheitlichen Theorie der Gamma- und der hypergeometrischen Funktionen. *Mathematische Annalen* 68 (1910), 305–337.
66. OBERHETTINGER, F. *Tables of Mellin Transforms*. Springer, 1974. 275 pages.

67. OBERHETTINGER, F. *Tables of Fourier Transforms and Fourier Transforms of Distributions*. Springer Verlag, 1990.
68. ODLYZKO, A. M. Enumeration of strings. In *Combinatorial Algorithms on Words* (1985), A. Apostolico and Z. Galil, Eds., vol. 12 of *NATO Advance Science Institute Series. Series F: Computer and Systems Sciences*, Springer Verlag, pp. 205–228.
69. PIPPENGER, N. An elementary approach to some analytic asymptotics. *SIAM Journal on Mathematical Analysis* 24, 5 (1993), 1361–1377.
70. PRODINGER, H. How to select a loser. *Discrete Mathematics* 120 (1993), 149–159.
71. PRODINGER, H. Hypothetical analyses: approximate counting in the style of Knuth, path length in the style of Flajolet. *Theoretical Computer Science* 100, 1 (1993), 243–251.
72. RÉGNIER, M. Evaluation des performances du hachage dynamique, 1983. Thèse de 3e cycle, Université Paris-Sud.
73. RÉGNIER, M. Analysis of grid file algorithms. *BIT* 25 (1985), 335–357.
74. REINGOLD, E. M., AND SUPOWIT, K. J. Probabilistic analysis of Divide-and-Conquer heuristics for minimum weighted Euclidean matching. *Networks*, 13 (1983), 49–66.
75. SEDGEWICK, R. Data movement in odd-even merging. *SIAM Journal on Computing* 7, 3 (1978), 239–272.
76. SEDGEWICK, R. *Algorithms*, second ed. Addison-Wesley, Reading, Mass., 1988.
77. SNEDDON, I. N. *The Use of Integral Transforms*. McGraw-Hill Inc., 1972.
78. STANLEY, R. P. *Enumerative Combinatorics*, vol. I. Wadsworth & Brooks/Cole, 1986.
79. SZPANKOWSKI, W. Some results on V-ary asymmetric tries. *Journal of Algorithms* 9 (1988), 224–244.
80. SZPANKOWSKI, W. Patricia tries again revisited. *Journal of the ACM* 37, 4 (1990), 691–711.
81. TITCHMARSH, E. C. *The Theory of Functions*, second ed. Oxford University Press, 1939.
82. TITCHMARSH, E. C. *Introduction to the Theory of Fourier Integrals*. Chelsea Publishing Company, 1986. Reprinted from the second edition, Oxford 1948.
83. TITCHMARSH, E. C., AND HEATH-BROWN, D. R. *The Theory of the Riemann Zeta-function*, second ed. Oxford Science Publications, 1986.
84. VITTER, J. S., AND FLAJOLET, P. Analysis of algorithms and data structures. In *Handbook of Theoretical Computer Science*, J. van Leeuwen, Ed., vol. A: Algorithms and Complexity. North Holland, 1990, ch. 9, pp. 431–524.
85. WHITTAKER, E. T., AND WATSON, G. N. *A Course of Modern Analysis*, fourth ed. Cambridge University Press, 1927. Reprinted 1973.
86. WIDDER, D. V. *The Laplace Transform*. Princeton University Press, 1941.
87. WILF, H. S. *Generatingfunctionology*. Academic Press, 1990.
88. WONG, R. *Asymptotic Approximations of Integrals*. Academic Press, 1989.

CONTENTS

Introduction	1
Principles.	
1. Basic properties	7
Functional properties.	
Inversion.	
2. The fundamental correspondence	12
Direct mapping.	
Converse mapping.	
Periodicities.	
3. Harmonic sums	20
4. Summatory formulæ	28
Euler-Maclaurin summation.	
Dyadic sums.	
Poisson's summation formula.	
Perron's formula.	
5. Catalan sums	33
6. The Bernoulli splitting process and dyadic sums	36
7. Other combinatorial examples	38
Reduction to standard harmonic sums.	
Nonstandard Dirichlet series.	
8. General conditions for Mellin asymptotics	41
Smallness of Mellin transforms.	
Growth of Dirichlet series.	
The transfer technique.	
9. Conclusions	49
References	51

Algorithms Project, INRIA, Rocquencourt, F-78153 Le Chesnay, France.

E-mail address: Philippe.Flajolet@inria.fr, Xavier.Gourdon@inria.fr, Philippe.Dumas@inria.fr



Unité de recherche INRIA Rocquencourt
Domaine de Voluceau - Rocquencourt - B.P. 105 - 78153 Le Chesnay Cedex (France)
Unité de recherche INRIA Lorraine - Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - B.P. 101 - 54602 Villers lès Nancy Cedex (France)
Unité de recherche INRIA Rennes - IRISA, Campus universitaire de Beaulieu 35042 Rennes Cedex (France)
Unité de recherche INRIA Rhône-Alpes 46, avenue Félix Viallet - 38031 Grenoble Cedex 1 (France)
Unité de recherche INRIA Sophia Antipolis - 2004, route des Lucioles - B.P. 93 - 06902 Sophia Antipolis Cedex (France)

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt - B.P. 105 - 78153 Le Chesnay Cedex (France)

ISSN 0249 - 6399



★ R R - 2 3 6 9 ★